

Odporúčania Európskej resuscitačnej rady pre kardiopulmonálnu resuscitáciu 2010

Sekcia 1 Súhrn

Resuscitation 2010; 81: 1219 - 1276

Zoznam literatúry

1. Nolan J. European Resuscitation Council Guidelines for resuscitation 2005. Section 1. Introduction. *Resuscitation* 2005;67(Suppl. 1):S3–6.
2. Nolan JP, Hazinski MF, Billi JE, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 1. Executive Summary. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.002, in press.
3. Nolan JP, Neumar RW, Adrie C, et al. Postcardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A Scientific Statement from the International Liaison Committee on Resuscitation; the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; the Council on Stroke. *Resuscitation* 2008;79:350–79.
4. Koster RW, Baubin MA, Caballero A, et al. European Resuscitation Council Guidelines for Resuscitation 2010. Section 2. Adult basic life support and use of automated external defibrillators. *Resuscitation* 2010;81:1277–92.
5. Deakin CD, Nolan JP, Sunde K, Koster RW. European Resuscitation Council Guidelines for Resuscitation 2010. Section 3. Electrical therapies: automated external defibrillators, defibrillation, cardioversion and pacing. *Resuscitation* 2010;81:1293–304.
6. Deakin CD, Nolan JP, Soar J, et al. European Resuscitation Council Guidelines for Resuscitation 2010. Section 4. Adult advanced life support. *Resuscitation* 2010;81:1305–52.
7. Arntz HR, Bossaert L, Danchin N, Nikolaou N. European Resuscitation Council Guidelines for Resuscitation 2010. Section 5. Initial management of acute coronary syndromes. *Resuscitation* 2010;81:1353–63.
8. Biarent D, Bingham R, Eich C, et al. European Resuscitation Council Guidelines for Resuscitation 2010. Section 6. Paediatric life support. *Resuscitation* 2010;81:1364–87.
9. Wyllie J, Richmond S. European Resuscitation Council Guidelines for Resuscitation 2010. Section 7. Resuscitation of babies at birth. *Resuscitation* 2010;81:1388–98.
10. Soar J, Perkins GD, Abbas G, et al. European Resuscitation Council Guidelines for Resuscitation 2010. Section 8. Cardiac arrest in special circumstances: electrolyte abnormalities, poisoning, drowning, accidental hypothermia, hyperthermia, asthma, anaphylaxis, cardiac surgery, trauma, pregnancy, electrocution. *Resuscitation* 2010;81:1399–431.
11. Soar J, Monsieurs KG, Ballance J, et al. European Resuscitation Council Guidelines for Resuscitation. Section 9. Principles of education in resuscitation. *Resuscitation* 2010;81:1432–42.
12. Lippert FK, Raffay V, Georgiou M, Steen PA, Bossaert L. European Resuscitation Council Guidelines for Resuscitation 2010. Section 10. The ethics of resuscitation and endoflife decisions. *Resuscitation* 2010;81:1443–9.
13. Koster RW, Sayre MR, Botha M, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 5. Adult basic life support. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.005, in press.
14. Sunde K, Jacobs I, Deakin CD, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 6. Defibrillation. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.025, in press.
15. Deakin CD, Morrison LJ, Morley PT, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 8. Advanced life support. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.027, in press.
16. Bossaert L, O'Connor RE, Arntz HR, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 9. Acute coronary syndromes. *Resuscitation*; doi:10.1016/j.resuscitation.2010.09.001, in press.
17. de Caen AR, Kleinman ME, Chameides L, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 10. Pediatric basic and advanced life support. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.028, in press.

18. Wyllie J, Perlman JM, Kattwinkel J, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 11. Neonatal resuscitation. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.029, in press.
19. Soar J, Mancini ME, Bhanji F, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 12. Education, implementation, and teams. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.030, in press.
20. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: global burden of disease study. *Lancet* 1997;349:1269–76.
21. Sans S, Kesteloot H, Kromhout D. The burden of cardiovascular diseases mortality in Europe. Task force of the European Society of Cardiology on cardiovascular mortality and morbidity statistics in Europe. *Eur Heart J* 1997;18:1231–48.
22. Zheng ZJ, Croft JB, Giles WH, Mensah GA. Sudden cardiac death in the United States, 1989 to 1998. *Circulation* 2001;104:2158–63.
- 22a. Atwood C, Eisenberg MS, Herlitz J, Rea TD. Incidence of EMStreated outofhospital cardiac arrest in Europe. *Resuscitation* 2005;67:75–80.
23. Nichol G, Thomas E, Callaway CW, et al. Regional variation in outofhospital cardiac arrest incidence and outcome. *JAMA* 2008;300:1423–31.
24. Hollenberg J, Herlitz J, Lindqvist J, et al. Improved survival after outofhospital cardiac arrest is associated with an increase in proportion of emergency crew—witnessed cases and bystander cardiopulmonary resuscitation. *Circulation* 2008;118:389–96.
25. Iwami T, Nichol G, Hiraide A, et al. Continuous improvements in “chain of survival” increased survival after outofhospital cardiac arrests: a largescale populationbased study. *Circulation* 2009;119:728–34.
26. Cobb LA, Fahrenbruch CE, Olsufka M, Copass MK. Changing incidence of outofhospital ventricular fibrillation, 1980–2000. *JAMA* 2002;288:3008–13.
27. Rea TD, Pearce RM, Raghunathan TE, et al. Incidence of outofhospital cardiac arrest. *Am J Cardiol* 2004;93:1455–60.
28. Vaillancourt C, Verma A, Trickett J, et al. Evaluating the effectiveness of dispatchassisted cardiopulmonary resuscitation instructions. *Acad Emerg Med* 2007;14:877–83.
29. Agarwal DA, Hess EP, Atkinson EJ, White RD. Ventricular fibrillation in Rochester, Minnesota: experience over 18 years. *Resuscitation* 2009;80:1253–8.
30. Ringh M, Herlitz J, Hollenberg J, Rosenqvist M, Svensson L. Out of hospital cardiac arrest outside home in Sweden, change in characteristics, outcome and availability for public access defibrillation. *Scand J Trauma Resusc Emerg Med* 2009;17:18.
31. Cummins R, Thies W. Automated external defibrillators and the Advanced Cardiac Life Support Program: a new initiative from the American Heart Association. *Am J Emerg Med* 1991;9:91–3.
32. Waalewijn RA, Nijpels MA, Tijssen JG, Koster RW. Prevention of deterioration of ventricular fibrillation by basic life support during outofhospital cardiac arrest. *Resuscitation* 2002;54:31–6.
33. Weisfeldt ML, Sitlani CM, Ornato JP, et al. Survival after application of automatic external defibrillators before arrival of the emergency medical system: evaluation in the resuscitation outcomes consortium population of 21 million. *J Am Coll Cardiol* 2010;55:1713–20.
34. van Alem AP, Vrenken RH, de Vos R, Tijssen JG, Koster RW. Use of automated external defibrillator by first responders in out of hospital cardiac arrest: prospective controlled trial. *BMJ* 2003;327:1312.
35. Sandroni C, Nolan J, Cavallaro F, Antonelli M. Inhospital cardiac arrest: incidence, prognosis and possible measures to improve survival. *Intensive Care Med* 2007;33:237–45.
36. Meaney PA, Nadkarni VM, Kern KB, Indik JH, Halperin HR, Berg RA. Rhythms and outcomes of adult inhospital cardiac arrest. *Crit Care Med* 2010;38:101–8.
37. Proceedings of the 2005 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care science with Treatment Recommendations. *Resuscitation* 2005;67:157–341.
38. International Liaison Committee on Resuscitation. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Circulation* 2005;112(Suppl. III):III1–136.
39. *J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 1265 39. Morley PT, Atkins DL, Billi JE, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 3. Evidence evaluation process. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.023, in press.
40. Billi JE, Zideman DA, Eigel B, Nolan JP, Montgomery WH, Nadkarni VM. Conflict of interest management before, during, and after the 2005 International Consensus Conference on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2005;67:171–3.

41. Shuster M, Billi JE, Bossaert L, et al. International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 4. Conflict of interest management before, during, and after the 2010 International Consensus Conference on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation*; doi:10.1016/j.resuscitation.2010.08.024, in press.
42. Valenzuela TD, Roe DJ, Cretin S, Spaite DW, Larsen MP. Estimating effectiveness of cardiac arrest interventions: a logistic regression survival model. *Circulation* 1997;96:3308–13.
43. Holmberg M, Holmberg S, Herlitz J. Factors modifying the effect of bystander cardiopulmonary resuscitation on survival in outofhospital cardiac arrest patients in Sweden. *Eur Heart J* 2001;22:511–9.
44. Holmberg M, Holmberg S, Herlitz J, Gardelov B. Survival after cardiac arrest outside hospital in Sweden. *Swedish Cardiac Arrest Registry. Resuscitation* 1998;36:29–36.
45. Waalewijn RA, Tijssen JG, Koster RW. Bystander initiated actions in outofhospital cardiopulmonary resuscitation: results from the Amsterdam Resuscitation Study (ARREST). *Resuscitation* 2001;50:273–9.
46. SOSKANTO Study Group. Cardiopulmonary resuscitation by bystanders with chest compression only (SOSKANTO): an observational study. *Lancet* 2007;369:920–6.
47. Iwami T, Kawamura T, Hiraide A, et al. Effectiveness of bystanderinitiated cardiaonly resuscitation for patients with outofhospital cardiac arrest. *Circulation* 2007;116:2900–7.
48. Weaver WD, Hill D, Fahrenbruch CE, et al. Use of the automatic external defibrillator in the management of outofhospital cardiac arrest. *N Engl J Med* 1988;319:661–6.
49. Auble TE, Menegazzi JJ, Paris PM. Effect of outofhospital defibrillation by basic life support providers on cardiac arrest mortality: a metaanalysis. *Ann Emerg Med* 1995;25:642–58.
50. Stiell IG, Wells GA, Field BJ, et al. Improved outofhospital cardiac arrest survival through the inexpensive optimization of an existing defibrillation program: OPALS study phase II. *Ontario Prehospital Advanced Life Support. JAMA* 1999;281:1175–81.
51. Stiell IG, Wells GA, DeMaio VJ, et al. Modifiable factors associated with improved cardiac arrest survival in a multicenter basic life support/ defibrillation system: OPALS Study Phase I results. *Ontario Prehospital Advanced Life Support. Ann Emerg Med* 1999;33:44–50.
52. Caffrey S. Feasibility of public access to defibrillation. *Curr Opin Crit Care* 2002;8:195–8.
53. O'Rourke MF, Donaldson E, Geddes JS. An airline cardiac arrest program. *Circulation* 1997;96:2849–53.
54. Page RL, Hamdan MH,McKenas DK. Defibrillation aboard a commercial aircraft. *Circulation* 1998;97:1429–30.
55. Valenzuela TD, Roe DJ, Nichol G, Clark LL, Spaite DW, Hardman RG. Outcomes of rapid defibrillation by security officers after cardiac arrest in casinos. *N Engl J Med* 2000;343:1206–9.
56. Waalewijn RA, de Vos R, Tijssen JG, Koster RW. Survival models for outofhospital cardiopulmonary resuscitation from the perspectives of the bystander, the first responder, and the paramedic. *Resuscitation* 2001;51:113–22.
57. Engdahl J, Abrahamsson P, Bang A, Lindqvist J, Karlsson T, Herlitz J. Is hospital care of major importance for outcome after outofhospital cardiac arrest? Experience acquired from patients with outofhospital cardiac arrest resuscitated by the same Emergency Medical Service and admitted to one of two hospitals over a 16year period in the municipality of Goteborg. *Resuscitation* 2000;43:201–11.
58. Langhelle A, Tyvold SS, Lexow K, Hapnes SA, Sunde K, Steen PA. Inhospital factors associated with improved outcome after outofhospital cardiac arrest. A comparison between four regions in Norway. *Resuscitation* 2003;56:247–63.
59. Carr BG, Goyal M, Band RA, et al. A national analysis of the relationship between hospital factors and postcardiac arrest mortality. *Intensive Care Med* 2009;35:505–11.
60. Liu JM, Yang Q, Pirrallo RG, Klein JP, Aufderheide TP. Hospital variability of outofhospital cardiac arrest survival. *Prehosp Emerg Care* 2008;12:339–46.
61. Carr BG, Kahn JM, Merchant RM, Kramer AA, Neumar RW. Interhospital variability in postcardiac arrest mortality. *Resuscitation* 2009;80:30–4.
62. Herlitz J, Engdahl J, Svensson L, Angquist KA, Silfverstolpe J, Holmberg S. Major differences in 1month survival between hospitals in Sweden among initial survivors of outofhospital cardiac arrest. *Resuscitation* 2006;70:404–9.
63. Keenan SP, Dodek P, Martin C, Priestap F, Norena M,WongH. Variation in length of intensive care unit stay after cardiac arrest: where you are is as important as who you are. *Crit Care Med* 2007;35:836–41.
64. Bahr J, Klingler H, Panzer W, Rode H, Kettler D. Skills of lay people in checking the carotid pulse. *Resuscitation* 1997;35:23–6.
65. Nyman J, Sihvonen M. Cardiopulmonary resuscitation skills in nurses and nursing students. *Resuscitation* 2000;47:179–84.

66. Tibballs J, Russell P. Reliability of pulse palpation by healthcare personnel to diagnose paediatric cardiac arrest. *Resuscitation* 2009;80:61–4.
67. Ruppert M, Reith MW, Widmann JH, et al. Checking for breathing: evaluation of the diagnostic capability of emergency medical services personnel, physicians, medical students, and medical laypersons. *Ann Emerg Med* 1999;34: 720–9.
68. Perkins GD, Stephenson B, Hulme J, Monsieurs KG. Birmingham assessment of breathing study (BABS). *Resuscitation* 2005;64:109–13.
69. Bobrow BJ, Zuercher M, Ewy GA, et al. Gasping during cardiac arrest in humans is frequent and associated with improved survival. *Circulation* 2008;118:2550–4.
70. Taylor RB, Brown CG, Bridges T, Werman HA, Ashton J, Hamlin RL. A model for regional blood flow measurements during cardiopulmonary resuscitation in a swine model. *Resuscitation* 1988;16:107–18.
71. Eftestol T, Sunde K, Steen PA. Effects of interrupting precordial compressions on the calculated probability of defibrillation success during outofhospital cardiac arrest. *Circulation* 2002;105:2270–3.
72. Aufderheide TP, Pirralo RG, Yannopoulos D, et al. Incomplete chest wall decompression: a clinical evaluation of CPR performance by EMS personnel and assessment of alternative manual chest compression–decompression techniques. *Resuscitation* 2005;64:353–62.
73. Yannopoulos D, McKnite S, Aufderheide TP, et al. Effects of incomplete chest wall decompression during cardiopulmonary resuscitation on coronary and cerebral perfusion pressures in a porcine model of cardiac arrest. *Resuscitation* 2005;64:363–72.
74. Ornato JP, Hallagan LF, McMahan SB, Peeples EH, Rostafinski AG. Attitudes of BCLS instructors about mouth-to-mouth resuscitation during the AIDS epidemic. *Ann Emerg Med* 1990;19:151–6.
75. Hew P, Brenner B, Kaufman J. Reluctance of paramedics and emergency medical technicians to perform mouth-to-mouth resuscitation. *J Emerg Med* 1997;15:279–84.
76. Chandra NC, Gruben KG, Tsitlik JE, et al. Observations of ventilation during resuscitation in a canine model. *Circulation* 1994;90:3070–5.
77. Kern KB, Hilwig RW, Berg RA, Sanders AB, Ewy GA. Importance of continuous chest compressions during cardiopulmonary resuscitation: improved outcome during a simulated single layrescuer scenario. *Circulation* 2002;105: 645–9.
78. Geddes LA, Rundell A, Otlewski M, Pargett M. How much lung ventilation is obtained with only chestcompression CPR? *Cardiovasc Eng* 2008;8:145–8.
79. Berg RA, Kern KB, Hilwig RW, et al. Assisted ventilation does not improve outcome in a porcine model of singlerescuer bystander cardiopulmonary resuscitation. *Circulation* 1997;95:1635–41.
80. Berg RA, Kern KB, Hilwig RW, Ewy GA. Assisted ventilation during ‘bystander’ CPR in a swine acute myocardial infarction model does not improve outcome. *Circulation* 1997;96:4364–71.
81. Turner I, Turner S, Armstrong V. Does the compression to ventilation ratio affect the quality of CPR: a simulation study. *Resuscitation* 2002;52:55–62.
82. Dorph E, WikL, Stromme TA, Eriksen M, Steen PA. Oxygen delivery and return of spontaneous circulation with ventilation:compression ratio 2:30 versus chest compressions only CPR in pigs. *Resuscitation* 2004;60:309–18.
83. Bohm K, Rosenqvist M, Herlitz J, Hollenberg J, Svensson L. Survival is similar after standard treatment and chest compression only in outofhospital bystander cardiopulmonary resuscitation. *Circulation* 2007;116:2908–12.
84. Kitamura T, Iwami T, Kawamura T, et al. Conventional and chestcompressiononly cardiopulmonary resuscitation by bystanders for children who have outofhospital cardiac arrests: a prospective, nationwide, populationbased cohort study. *Lancet* 2010.
85. Kitamura T, Iwami T, Kawamura T, Nagao K, Tanaka H, Hiraide A. Bystanderinitiated rescue breathing for outofhospital cardiac arrests of noncardiac origin. *Circulation* 2010;122:293–9.
86. Peberdy MA, Ottingham LV, Groh WJ, et al. Adverse events associated with lay emergency response programs: the public access defibrillation trial experience. *Resuscitation* 2006;70:59–65.
87. Sugerma NT, Edelson DP, Leary M, et al. Rescuer fatigue during actual inhospital cardiopulmonary resuscitation with audiovisual feedback: a prospective multicenter study. *Resuscitation* 2009;80:981–4.
88. Hallstrom AP, Ornato JP, Weisfeldt M, et al. Publicaccess defibrillation and survival after outofhospital cardiac arrest. *N Engl J Med* 2004;351:637–46.
89. Hoke RS, Heinroth K, Trappe HJ, Werdan K. Is external defibrillation an electric threat for bystanders? *Resuscitation* 2009;80:395–401.
90. Dickinson CL, Hall CR, Soar J. Accidental shock to rescuer during successful defibrillation of ventricular fibrillation—a case of human involuntary automaticity. *Resuscitation* 2008;76:489.
91. Cydulka RK, Connor PJ, Myers TF, Pavza G, Parker M. Prevention of oral bacterial flora transmission by using mouth-to-mask ventilation during CPR. *J Emerg Med* 1991;9:317–21.

92. Blenkarn JI, Buckingham SE, Zideman DA. Prevention of transmission of infection during mouth-to-mouth resuscitation. *Resuscitation* 1990;19:151–7.
93. Turner S, Turner I, Chapman D, et al. A comparative study of the 1992 and 1997 recovery positions for use in the UK. *Resuscitation* 1998;39:153–60.
94. Handley AJ. Recovery position. *Resuscitation* 1993;26:93–5.
95. Anon. Guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care—an international consensus on science. *Resuscitation* 2000;46:1–447.
96. Fingerhut LA, Cox CS, Warner M. International comparative analysis of injury mortality. Findings from the ICE on injury statistics. International Collaborative Effort on Injury Statistics. *Adv Data* 1998:1–20.
- 1266 J.P. Nolan et al. / *Resuscitation* 81 (2010) 1219–1276
97. White RD, Bunch TJ, Hankins DG. Evolution of a communitywide early defibrillation programme experience over 13 years using police/fire personnel and paramedics as responders. *Resuscitation* 2005;65:279–83.
98. Mosesso Jr VN, Davis EA, Auble TE, Paris PM, Yealy DM. Use of automated external defibrillators by police officers for treatment of out-of-hospital cardiac arrest. *Ann Emerg Med* 1998;32:200–7.
99. The Public Access Defibrillation Trial Investigators. Public access defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:637–46.
100. Kitamura T, Iwami T, Kawamura T, Nagao K, Tanaka H, Hiraide A. Nationwide public access defibrillation in Japan. *N Engl J Med* 2010;362:994–1004.
101. Bardy GH, Lee KL, Mark DB, et al. Home use of automated external defibrillators for sudden cardiac arrest. *N Engl J Med* 2008;358:1793–804.
102. Zafari AM, Zarter SK, Heggen V, et al. A program encouraging early defibrillation results in improved in-hospital resuscitation efficacy. *J Am Coll Cardiol* 2004;44:846–52.
103. Destro A, Marzaloni M, Sermasi S, Rossi F. Automatic external defibrillators in the hospital as well? *Resuscitation* 1996;31:39–43.
104. Spearpoint KG, Gruber PC, Brett SJ. Impact of the Immediate Life Support course on the incidence and outcome of in-hospital cardiac arrest calls: an observational study over 6 years. *Resuscitation* 2009;80:638–43.
105. Cummins RO, Eisenberg MS, Litwin PE, Graves JR, Hearne TR, Hallstrom AP. Automatic external defibrillators used by emergency medical technicians: a controlled clinical trial. *JAMA* 1987;257:1605–10.
106. Stults KR, Brown DD, Kerber RE. Efficacy of an automated external defibrillator in the management of out-of-hospital cardiac arrest: validation of the diagnostic algorithm and initial clinical experience in a rural environment. *Circulation* 1986;73:701–9.
107. Kramer-Johansen J, Edelson DP, Abella BS, Becker LB, Wik L, Steen PA. Pauses in chest compression and inappropriate shocks: a comparison of manual and semiautomatic defibrillation attempts. *Resuscitation* 2007;73:212–20.
108. Pytte M, Pedersen TE, Ottem J, Rokvam AS, Sunde K. Comparison of hands-off time during CPR with manual and semiautomatic defibrillation in a manikin model. *Resuscitation* 2007;73:131–6.
109. Forcina MS, Farhat AY, O’Neil WW, Haines DE. Cardiac arrest survival after implementation of automated external defibrillator technology in the in-hospital setting. *Crit Care Med* 2009;37:1229–36.
110. Edelson DP, Abella BS, Kramer-Johansen J, et al. Effects of compression depth and pre-shock pauses predict defibrillation failure during cardiac arrest. *Resuscitation* 2006;71:137–45.
111. Yu T, Weil MH, Tang W, et al. Adverse outcomes of interrupted precordial compression during automated defibrillation. *Circulation* 2002;106:368–72.
112. Gundersen K, Kvaloy JT, Kramer-Johansen J, Steen PA, Eftestol T. Development of the probability of return of spontaneous circulation in intervals without chest compressions during out-of-hospital cardiac arrest: an observational study. *BMC Med* 2009;7:6.
113. Lloyd MS, Heeke B, Walter PF, Langberg JJ. Hands-on defibrillation: an analysis of electrical current flow through rescuers in direct contact with patients during biphasic external defibrillation. *Circulation* 2008;117:2510–4.
114. Bojar RM, Payne DD, Rastegar H, Diehl JT, Cleveland RJ. Use of self-adhesive external defibrillator pads for complex cardiac surgical procedures. *Ann Thorac Surg* 1988;46:587–8.
115. Bradbury N, Hyde D, Nolan J. Reliability of ECG monitoring with a gel pad/paddle combination after defibrillation. *Resuscitation* 2000;44:203–6.
116. Brown J, Rogers J, Soar J. Cardiac arrest during surgery and ventilation in the prone position: a case report and systematic review. *Resuscitation* 2001;50:233–8.
117. Perkins GD, Davies RP, Soar J, Thickett DR. The impact of manual defibrillation technique on no-flow time during simulated cardiopulmonary resuscitation. *Resuscitation* 2007;73:109–14.
118. Wilson RF, Sirna S, White CW, Kerber RE. Defibrillation of high-risk patients during coronary angiography using self-adhesive, preapplied electrode pads. *Am J Cardiol* 1987;60:380–2.

119. Stults KR, Brown DD, Cooley F, Kerber RE. Selfadhesive monitor/defibrillation pads improve prehospital defibrillation success. *Ann Emerg Med* 1987;16:872–7.
120. Callaway CW, Sherman LD, Mosesso Jr VN, Dietrich TJ, Holt E, Clarkson MC. Scaling exponent predicts defibrillation success for outofhospital ventricular fibrillation cardiac arrest. *Circulation* 2001;103:1656–61.
121. Eftestol T, Sunde K, Aase SO, Husoy JH, Steen PA. Predicting outcome of defibrillation by spectral characterization and nonparametric classification of ventricular fibrillation in patients with outofhospital cardiac arrest. *Circulation* 2000;102:1523–9.
122. Eftestol T, Wik L, Sunde K, Steen PA. Effects of cardiopulmonary resuscitation on predictors of ventricular fibrillation defibrillation success during outofhospital cardiac arrest. *Circulation* 2004;110:10–5.
123. Weaver WD, Cobb LA, Dennis D, Ray R, Hallstrom AP, Copass MK. Amplitude of ventricular fibrillation waveform and outcome after cardiac arrest. *Ann Intern Med* 1985;102:53–5.
124. Brown CG, Dzwonczyk R. Signal analysis of the human electrocardiogram during ventricular fibrillation: frequency and amplitude parameters as predictors of successful countershock. *Ann Emerg Med* 1996;27:184–8.
125. Callahan M, Braun O, Valentine W, Clark DM, Zegans C. Prehospital cardiac arrest treated by urban firstresponders: profile of patient response and prediction of outcome by ventricular fibrillation waveform. *Ann Emerg Med* 1993;22:1664–77.
126. Strohmer HU, Lindner KH, Brown CG. Analysis of the ventricular fibrillation ECG signal amplitude and frequency parameters as predictors of countershock success in humans. *Chest* 1997;111:584–9.
127. Strohmer HU, Eftestol T, Sunde K, et al. The predictive value of ventricular fibrillation electrocardiogram signal frequency and amplitude variables in patients with outofhospital cardiac arrest. *Anesth Analg* 2001;93:1428–33.
128. Podbregar M, Kovacic M, Podbregar Mars A, Brezocnik M. Predicting defibrillation success by ‘genetic’ programming in patients with outofhospital cardiac arrest. *Resuscitation* 2003;57:153–9.
129. Menegazzi JJ, Callaway CW, Sherman LD, et al. Ventricular fibrillation scaling exponent can guide timing of defibrillation and other therapies. *Circulation* 2004;109:926–31.
130. Povoas HP, Weil MH, Tang W, Bisera J, Klouche K, Barbatsis A. Predicting the success of defibrillation by electrocardiographic analysis. *Resuscitation* 2002;53:77–82.
131. Noc M, Weil MH, Tang W, Sun S, Pernat A, Bisera J. Electrocardiographic prediction of the success of cardiac resuscitation. *Crit Care Med* 1999;27:708–14.
132. Strohmer HU, Lindner KH, Keller A, Lindner IM, Pfenninger EG. Spectral analysis of ventricular fibrillation and closedchest cardiopulmonary resuscitation. *Resuscitation* 1996;33:155–61.
133. Noc M, Weil MH, Gazmuri RJ, Sun S, Biscera J, Tang W. Ventricular fibrillation voltage as a monitor of the effectiveness of cardiopulmonary resuscitation. *J Lab Clin Med* 1994;124:421–6.
134. Lightfoot CB, Nremt P, Callaway CW, et al. Dynamic nature of electrocardiographic waveform predicts rescue shock outcome in porcine ventricular fibrillation. *Ann Emerg Med* 2003;42:230–41.
135. Marn Pernat A, Weil MH, Tang W, Pernat A, Bisera J. Optimizing timing of ventricular defibrillation. *Crit Care Med* 2001;29:2360–5.
136. Hamprecht FA, Achleitner U, Krismer AC, et al. Fibrillation power, an alternative method of ECG spectral analysis for prediction of countershock success in a porcine model of ventricular fibrillation. *Resuscitation* 2001;50:287–96.
137. Amann A, Achleitner U, Antretter H, et al. Analysing ventricular fibrillation ECG signals and predicting defibrillation success during cardiopulmonary resuscitation employing N(alpha) histograms. *Resuscitation* 2001;50:77–85.
138. Brown CG, Griffith RF, Van Ligten P, et al. Median frequency—a new parameter for predicting defibrillation success rate. *Ann Emerg Med* 1991;20:787–9.
139. Amann A, Rheinberger K, Achleitner U, et al. The prediction of defibrillation outcome using a new combination of mean frequency and amplitude in porcine models of cardiac arrest. *Anesth Analg* 2002;95:716–22 [table of contents].
140. Deakin CD, Nolan JP. European Resuscitation Council guidelines for resuscitation 2005. Section 3. Electrical therapies: automated external defibrillators, defibrillation, cardioversion and pacing. *Resuscitation* 2005;67(Suppl. 1):S25–37.
141. Cobb LA, Fahrenbruch CE, Walsh TR, et al. Influence of cardiopulmonary resuscitation prior to defibrillation in patients with outofhospital ventricular fibrillation. *JAMA* 1999;281:1182–8.
142. Wik L, Hansen TB, Fylling F, et al. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with outofhospital ventricular fibrillation: a randomized trial. *JAMA* 2003;289:1389–95.

143. Baker PW, Conway J, Cotton C, et al. Defibrillation or cardiopulmonary resuscitation first for patients with outofhospital cardiac arrests found by paramedics to be in ventricular fibrillation? A randomised control trial. *Resuscitation* 2008;79:424–31.
144. Jacobs IG, Finn JC, Oxer HF, Jelinek GA. CPR before defibrillation in outofhospital cardiac arrest: a randomized trial. *Emerg Med Australas* 2005;17:39–45.
145. Hayakawa M, Gando S, Okamoto H, Asai Y, Uegaki S, Makise H. Shortening of cardiopulmonary resuscitation time before the defibrillation worsens the outcome in outofhospital VF patients. *Am J Emerg Med* 2009;27:470–4.
146. Bradley SM, Gabriel EE, Aufderheide TP, et al. Survival Increases with CPR by Emergency Medical Services before defibrillation of outofhospital ventricular fibrillation or ventricular tachycardia: observations from the resuscitation outcomes consortium. *Resuscitation* 2010;81:155–62.
147. Christenson J, Andrusiek D, EversonStewart S, et al. Chest compression fraction determines survival in patients with outofhospital ventricular fibrillation. *Circulation* 2009;120:1241–7.
148. Olasveengen TM, Vik E, Kuzovlev A, Sunde K. Effect of implementation of new resuscitation guidelines on quality of cardiopulmonary resuscitation and survival. *Resuscitation* 2009;80:407–11.
149. Bobrow BJ, Clark LL, Ewy GA, et al. Minimally interrupted cardiac resuscitation by emergency medical services for outofhospital cardiac arrest. *JAMA* 2008;299:1158–65.
150. Rea TD, Helbock M, Perry S, et al. Increasing use of cardiopulmonary resuscitation during outofhospital ventricular fibrillation arrest: survival implications of guideline changes. *Circulation* 2006;114:2760–5.
151. Steinmetz J, Barnung S, Nielsen SL, Risom M, Rasmussen LS. Improved survival after an outofhospital cardiac arrest using new guidelines. *Acta Anaesthesiol Scand* 2008;52:908–13.
152. Jost D, Degrange H, Verret C, et al. DEFI 2005: a randomized controlled trial of the effect of automated external defibrillator cardiopulmonary resuscitation protocol on outcome from outofhospital cardiac arrest. *Circulation* 2010;121:1614–22.
153. van Alem AP, Chapman FW, Lank P, Hart AA, Koster RW. A prospective, randomised and blinded comparison of first shock success of monophasic and biphasic waveforms in outofhospital cardiac arrest. *Resuscitation* 2003;58:17–24.
- J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 1267
154. Carpenter J, Rea TD, Murray JA, Kudenchuk PJ, Eisenberg MS. Defibrillation waveform and postshock rhythm in outofhospital ventricular fibrillation cardiac arrest. *Resuscitation* 2003;59:189–96.
155. Morrison LJ, Dorian P, Long J, et al. Outofhospital cardiac arrest rectilinear biphasic to monophasic damped sine defibrillation waveforms with advanced life support intervention trial (ORBIT). *Resuscitation* 2005;66:149–57.
156. Mittal S, Ayati S, Stein KM, et al. Transthoracic cardioversion of atrial fibrillation: comparison of rectilinear biphasic versus damped sine wave monophasic shocks. *Circulation* 2000;101:1282–7.
157. Page RL, Kerber RE, Russell JK, et al. Biphasic versus monophasic shock waveform for conversion of atrial fibrillation: the results of an international randomized, doubleblind multicenter trial. *J Am Coll Cardiol* 2002;39:1956–63.
158. Koster RW, Dorian P, Chapman FW, Schmitt PW, O’Grady SG, Walker RG. A randomized trial comparing monophasic and biphasic waveform shocks for external cardioversion of atrial fibrillation. *Am Heart J* 2004;147:e20.
159. Ambler JJ, Deakin CD. A randomized controlled trial of efficacy and ST change following use of the WelchAllyn MRL PIC biphasic waveform versus damped sine monophasic waveform for external DC cardioversion. *Resuscitation* 2006;71:146–51.
160. Martens PR, Russell JK, Wolcke B, et al. Optimal response to cardiac arrest study: defibrillation waveform effects. *Resuscitation* 2001;49:233–43.
161. Gliner BE, Jorgenson DB, Poole JE, et al. Treatment of outofhospital cardiac arrest with a lowenergy impedancecompensating biphasic waveform automatic external defibrillator. *The LIFE Investigators. Biomed Instrum Technol* 1998;32:631–44.
162. White RD, Blackwell TH, Russell JK, Snyder DE, Jorgenson DB. Transthoracic impedance does not affect defibrillation, resuscitation or survival in patients with outofhospital cardiac arrest treated with a nonescalating biphasic waveform defibrillator. *Resuscitation* 2005;64:63–9.
163. Stiell IG, Walker RG, Nesbitt LP, et al. BIPHASIC trial: a randomized comparison of fixed lower versus escalating higher energy levels for defibrillation in outofhospital cardiac arrest. *Circulation* 2007;115:1511–7.
164. Walsh SJ, McClelland AJ, Owens CG, et al. Efficacy of distinct energy delivery protocols comparing two biphasic defibrillators for cardiac arrest. *Am J Cardiol* 2004;94:378–80.

165. Higgins SL, Herre JM, Epstein AE, et al. A comparison of biphasic and monophasic shocks for external defibrillation. *Physiocontrol biphasic investigators. Prehosp Emerg Care* 2000;4:305–13.
166. Berg RA, Samson RA, Berg MD, et al. Better outcome after pediatric defibrillation dosage than adult dosage in a swine model of pediatric ventricular fibrillation. *J Am Coll Cardiol* 2005;45:786–9.
167. Killingsworth CR, Melnick SB, Chapman FW, et al. Defibrillation threshold and cardiac responses using an external biphasic defibrillator with pediatric and adult adhesive patches in pediatric-sized piglets. *Resuscitation* 2002;55:177–85.
168. Tang W, Weil MH, Sun S, et al. The effects of biphasic waveform design on postresuscitation myocardial function. *J Am Coll Cardiol* 2004;43:1228–35.
169. Xie J, Weil MH, Sun S, et al. High-energy defibrillation increases the severity of postresuscitation myocardial dysfunction. *Circulation* 1997;96:683–8.
170. Lown B. Electrical reversion of cardiac arrhythmias. *Br Heart J* 1967;29:469–89.
171. Boodhoo L, Mitchell AR, Bordoli G, Lloyd G, Patel N, Sulke N. DC cardioversion of persistent atrial fibrillation: a comparison of two protocols. *Int J Cardiol* 2007;114:16–21.
172. Boos C, Thomas MD, Jones A, Clarke E, Wilbourne G, More RS. Higher energy monophasic DC cardioversion for persistent atrial fibrillation: is it time to start at 360 joules? *Ann Noninvasive Electrocardiol* 2003;8:121–6.
173. Glover BM, Walsh SJ, McCann CJ, et al. Biphasic energy selection for transthoracic cardioversion of atrial fibrillation. The BEST AF Trial. *Heart* 2008;94:884–7.
174. Rashba EJ, Gold MR, Crawford FA, Leman RB, Peters RW, Shorofsky SR. Efficacy of transthoracic cardioversion of atrial fibrillation using a biphasic, truncated exponential shock waveform at variable initial shock energies. *Am J Cardiol* 2004;94:1572–4.
175. Pinski SL, Sgarbossa EB, Ching E, Trohman RG. A comparison of 50J versus 100J shocks for direct-current cardioversion of atrial flutter. *Am Heart J* 1999;137:439–42.
176. Alatawi F, Gurevitz O, White R. Prospective, randomized comparison of two biphasic waveforms for the efficacy and safety of transthoracic biphasic cardioversion of atrial fibrillation. *Heart Rhythm* 2005;2:382–7.
177. Kerber RE, Martins JB, Kienzle MG, et al. Energy, current, and success in defibrillation and cardioversion: clinical studies using an automated impedance-based method of energy adjustment. *Circulation* 1988;77:1038–46.
178. Kerber RE, Kienzle MG, Olshansky B, et al. Ventricular tachycardia rate and morphology determine energy and current requirements for transthoracic cardioversion. *Circulation* 1992;85:158–63.
179. Stockwell B, Bellis G, Morton G, et al. Electrical injury during “hands on” defibrillation—a potential risk of internal cardioverter defibrillators? *Resuscitation* 2009;80:832–4.
180. Nolan J, Soar J, Eikeland H. The chain of survival. *Resuscitation* 2006;71: 270–1.
181. Gwinnutt CL, Columb M, Harris R. Outcome after cardiac arrest in adults in UK hospitals: effect of the 1997 guidelines. *Resuscitation* 2000;47:125–35.
182. Peberdy MA, Kaye W, Ornato JP, et al. Cardiopulmonary resuscitation of adults in the hospital: a report of 14720 cardiac arrests from the National Registry of Cardiopulmonary Resuscitation. *Resuscitation* 2003;58:297–308.
183. Smith GB. In-hospital cardiac arrest: is it time for an in-hospital ‘chain of prevention’? *Resuscitation* 2010.
184. National Confidential Enquiry into Patient Outcome and Death. An acute problem? London: NCEPOD; 2005.
185. Hodgetts TJ, Kenward G, Vlackonikolis I, et al. Incidence, location and reasons for avoidable in-hospital cardiac arrest in a district general hospital. *Resuscitation* 2002;54:115–23.
186. Kaue J, Smith G, Prytherch D, Parr M, Flabouris A, Hillman K. A comparison of antecedents to cardiac arrests, deaths and emergency intensive care admissions in Australia and New Zealand, and the United Kingdom—the ACADEMIA study. *Resuscitation* 2004;62:275–82.
187. Castagna J, Weil MH, Shubin H. Factors determining survival in patients with cardiac arrest. *Chest* 1974;65:527–9.
188. Herlitz J, Bang A, Aune S, Ekstrom L, Lundstrom G, Holmberg S. Characteristics and outcome among patients suffering in-hospital cardiac arrest in monitored and nonmonitored areas. *Resuscitation* 2001;48:125–35.
189. Campello G, Granja C, Carvalho F, Dias C, Azevedo LF, Costa-Pereira A. Immediate and long-term impact of medical emergency teams on cardiac arrest prevalence and mortality: a plea for periodic basic life support training programs. *Crit Care Med* 2009;37:3054–61.
190. Bellomo R, Goldsmith D, Uchino S, et al. A prospective before-and-after trial of a medical emergency team. *Med J Aust* 2003;179:283–7.

191. Bellomo R, Goldsmith D, Uchino S, et al. Prospective controlled trial of effect of medical emergency team on postoperative morbidity and mortality rates. *Crit Care Med* 2004;32:916–21.
192. DeVita MA, Smith GB, Adam SK, et al. “Identifying the hospitalised patient in crisis”—a consensus conference on the afferent limb of rapid response systems. *Resuscitation* 2010;81:375–82.
193. Goldhill DR, Worthington L, Mulcahy A, Tarling M, Sumner A. The patientatrisk team: identifying and managing seriously ill ward patients. *Anaesthesia* 1999;54:853–60.
194. Hodgetts TJ, Kenward G, Vlachonikolis IG, Payne S, Castle N. The identification of risk factors for cardiac arrest and formulation of activation criteria to alert a medical emergency team. *Resuscitation* 2002;54:125–31.
195. Subbe CP, Davies RG, Williams E, Rutherford P, Gemmell L. Effect of introducing the modified early warning score on clinical outcomes, cardiopulmonary arrests and intensive care utilisation in acute medical admissions. *Anaesthesia* 2003;58:797–802.
196. Armitage M, Eddleston J, Stokes T. Recognising and responding to acute illness in adults in hospital: summary of NICE guidance. *BMJ* 2007;335:258–9.
197. Hillman K, Chen J, Cretikos M, et al. Introduction of the medical emergency team (MET) system: a clusterrandomised controlled trial. *Lancet* 2005;365:2091–7.
198. Lee A, Bishop G, Hillman KM, Daffurn K. The Medical Emergency Team. *Anaesth Intensive Care* 1995;23:183–6.
199. Devita MA, Bellomo R, Hillman K, et al. Findings of the first consensus conference on medical emergency teams. *Crit Care Med* 2006;34:2463–78.
200. Ball C, Kirkby M, Williams S. Effect of the critical care outreach team on patient survival to discharge from hospital and readmission to critical care: nonrandomised population based study. *BMJ* 2003;327:1014.
201. Critical care outreach 2003: progress in developing services. The National Outreach Report. London, UK: Department of Health and National Health Service Modernisation Agency; 2003.
202. Chan PS, Jain R, Nallmothu BK, Berg RA, Sasson C. Rapid response teams: a systematic review and metaanalysis. *Arch Intern Med* 2010;170:18–26.
203. Parr MJ, Hadfield JH, Flabouris A, Bishop G, Hillman K. The Medical Emergency Team: 12 month analysis of reasons for activation, immediate outcome and notforresuscitation orders. *Resuscitation* 2001;50:39–44.
204. Smith GB. Increased do not attempt resuscitation decision making in hospitals with a medical emergency teams system—cause and effect? *Resuscitation* 2008;79:346–7.
205. Chen J, Flabouris A, Bellomo R, Hillman K, Finfer S. The Medical Emergency Team System and notforresuscitation orders: results from the MERIT study. *Resuscitation* 2008;79:391–7.
206. Jones DA, McIntyre T, Baldwin I, Mercer I, Kattula A, Bellomo R. The medical emergency team and endoflife care: a pilot study. *Crit Care Resusc* 2007;9:151–6.
207. Excellence NifHaC. NICE clinical guideline 50 acutely ill patients in hospital: recognition of and response to acute illness in adults in hospital. London: National Institute for Health and Clinical Excellence; 2007.
208. Marshall S, Harrison J, Flanagan B. The teaching of a structured tool improves the clarity and content of interprofessional clinical communication. *Qual Saf Health Care* 2009;18:137–40.
209. Muller D, Agrawal R, Arntz HR. How sudden is sudden cardiac death? *Circulation* 2006;114:1146–50.
210. Amital H, Glikson M, Burstein M, et al. Clinical characteristics of unexpected death among young enlisted military personnel: results of a threedecade retrospective surveillance. *Chest* 2004;126:528–33.
211. Basso C, Maron BJ, Corrado D, Thiene G. Clinical profile of congenital coronary artery anomalies with origin from the wrong aortic sinus leading to sudden death in young competitive athletes. *J Am Coll Cardiol* 2000;35:1493–501.
212. Corrado D, Basso C, Thiene G. Sudden cardiac death in young people with apparently normal heart. *Cardiovasc Res* 2001;50:399–408.
213. Drory Y, Turetz Y, Hiss Y, et al. Sudden unexpected death in persons less than 40 years of age. *Am J Cardiol* 1991;68:1388–92.
214. Kramer MR, Drori Y, Lev B. Sudden death in young soldiers. High incidence of syncope prior to death. *Chest* 1988;93:345–7.
- 1268 *J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 215. Quigley F, Greene M, O’Connor D, Kelly F. A survey of the causes of sudden cardiac death in the under 35yearage group. *Ir Med J* 2005;98:232–5.
216. Wisten A, Forsberg H, Krantz P, Messner T. Sudden cardiac death in 15–35year olds in Sweden during 1992–99. *J Intern Med* 2002;252:529–36.

217. Wisten A, Messner T. Young Swedish patients with sudden cardiac death have a lifestyle very similar to a control population. *Scand Cardiovasc J* 2005;39:137–42.
218. Wisten A, Messner T. Symptoms preceding sudden cardiac death in the young are common but often misinterpreted. *Scand Cardiovasc J* 2005;39: 143–9.
219. Morrison LJ, Visentin LM, Kiss A, et al. Validation of a rule for termination of resuscitation in outofhospital cardiac arrest. *N Engl J Med* 2006;355: 478–87.
220. Gabbott D, Smith G, Mitchell S, et al. Cardiopulmonary resuscitation standards for clinical practice and training in the UK. *Resuscitation* 2005;64:13–9.
221. Dyson E, Smith GB. Common faults in resuscitation equipment—guidelines for checking equipment and drugs used in adult cardiopulmonary resuscitation. *Resuscitation* 2002;55:137–49.
222. Perkins GD, Roberts C, Gao F. Delays in defibrillation: influence of different monitoring techniques. *Br J Anaesth* 2002;89:405–8.
223. Featherstone P, Chalmers T, Smith GB. RSVP: a system for communication of deterioration in hospital patients. *Br J Nurs* 2008;17:860–4.
224. Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during inhospital cardiac arrest. *JAMA* 2005;293:305–10.
225. Abella BS, Sandbo N, Vassilatos P, et al. Chest compression rates during cardiopulmonary resuscitation are suboptimal: a prospective study during inhospital cardiac arrest. *Circulation* 2005;111:428–34.
226. Stiell IG, Wells GA, Field B, et al. Advanced cardiac life support in outofhospital cardiac arrest. *N Engl J Med* 2004;351:647–56.
227. Olasveengen TM, Sunde K, Brunborg C, Thowsen J, Steen PA, Wik L. Intravenous drug administration during outofhospital cardiac arrest: a randomized trial. *JAMA* 2009;302:2222–9.
228. Herlitz J, Ekstrom L, Wennerblom B, Axelsson A, Bang A, Holmberg S. Adrenaline in outofhospital ventricular fibrillation. Does it make any difference? *Resuscitation* 1995;29:195–201.
229. Holmberg M, Holmberg S, Herlitz J. Low chance of survival among patients requiring adrenaline (epinephrine) or intubation after outofhospital cardiac arrest in Sweden. *Resuscitation* 2002;54:37–45.
230. Sunde K, Eftestol T, Askenberg C, Steen PA. Quality assessment of defibrillation and advanced life support using data from the medical control module of the defibrillator. *Resuscitation* 1999;41:237–47.
231. Rea TD, Shah S, Kudenchuk PJ, Copass MK, Cobb LA. Automated external defibrillators: to what extent does the algorithm delay CPR? *Ann Emerg Med* 2005;46:132–41.
232. van Alem AP, Sanou BT, Koster RW. Interruption of cardiopulmonary resuscitation with the use of the automated external defibrillator in outofhospital cardiac arrest. *Ann Emerg Med* 2003;42:449–57.
233. Pytte M, Kramer-Johansen J, Eilevstjonn J, et al. Haemodynamic effects of adrenaline (epinephrine) depend on chest compression quality during cardiopulmonary resuscitation in pigs. *Resuscitation* 2006;71:369–78.
234. Prengel AW, Lindner KH, Ensinger H, Grunert A. Plasma catecholamine concentrations after successful resuscitation in patients. *Crit Care Med* 1992;20:609–14.
235. Bhende MS, Thompson AE. Evaluation of an endtidal CO₂ detector during pediatric cardiopulmonary resuscitation. *Pediatrics* 1995;95:395–9.
236. Sehra R, Underwood K, Checchia P. End tidal CO₂ is a quantitative measure of cardiac arrest. *Pacing Clin Electrophysiol* 2003;26:515–7.
237. Amir O, Schliamser JE, Nemer S, Arie M. Ineffectiveness of precordial thump for cardioversion of malignant ventricular tachyarrhythmias. *Pacing Clin Electrophysiol* 2007;30:153–6.
238. Haman L, Parizek P, Vojacek J. Precordial thump efficacy in termination of induced ventricular arrhythmias. *Resuscitation* 2009;80:14–6.
239. Pellis T, Kette F, Lovisa D, et al. Utility of precordial thump for treatment of out of hospital cardiac arrest: a prospective study. *Resuscitation* 2009;80:17–23.
240. Kohl P, King AM, Boulin C. Antiarrhythmic effects of acute mechanical stimulation. In: Kohl P, Sachs F, Franz MR, editors. *Cardiac mechanoelectric feedback and arrhythmias: from pipette to patient*. Philadelphia: Elsevier Saunders; 2005. p. 304–14.
241. Caldwell G, Millar G, Quinn E, Vincent R, Chamberlain DA. Simple mechanical methods for cardioversion: defence of the precordial thump and cough version. *BMJ (Clin Res Ed)* 1985;291:627–30.
242. Wenzel V, Lindner KH, Augenstein S, et al. Intraosseous vasopressin improves coronary perfusion pressure rapidly during cardiopulmonary resuscitation in pigs. *Crit Care Med* 1999;27:1565–9.
243. Shavit I, Hoffmann Y, Galbraith R, Waisman Y. Comparison of two mechanical intraosseous infusion devices: a pilot, randomized crossover trial. *Resuscitation* 2009;80:1029–33.
244. Kudenchuk PJ, Cobb LA, Copass MK, et al. Amiodarone for resuscitation after outofhospital cardiac arrest due to ventricular fibrillation. *N Engl J Med* 1999;341:871–8.

245. Dorian P, Cass D, Schwartz B, Cooper R, Gelaznikas R, Barr A. Amiodarone as compared with lidocaine for shockresistant ventricular fibrillation. *N Engl J Med* 2002;346:884–90.
246. Thel MC, Armstrong AL, McNulty SE, Califf RM, O'Connor CM. Randomised trial of magnesium in inhospital cardiac arrest. Duke Internal Medicine Housestaff. *Lancet* 1997;350:1272–6.
247. Allegra J, Lavery R, Cody R, et al. Magnesium sulfate in the treatment of refractory ventricular fibrillation in the prehospital setting. *Resuscitation* 2001;49:245–9.
248. Fatovich D, Prentice D, Dobb G. Magnesium in inhospital cardiac arrest. *Lancet* 1998;351:446.
249. Hassan TB, Jagger C, Barnett DB. A randomised trial to investigate the efficacy of magnesium sulphate for refractory ventricular fibrillation. *Emerg Med J* 2002;19:57–62.
250. Miller B, Craddock L, Hoffenberg S, et al. Pilot study of intravenous magnesium sulfate in refractory cardiac arrest: safety data and recommendations for future studies. *Resuscitation* 1995;30:3–14.
251. Stiell IG, Wells GA, Hebert PC, Laupacis A, Weitzman BN. Association of drug therapy with survival in cardiac arrest: limited role of advanced cardiac life support drugs. *Acad Emerg Med* 1995;2:264–73.
252. Engdahl J, Bang A, Lindqvist J, Herlitz J. Can we define patients with no and those with some chance of survival when found in asystole out of hospital? *Am J Cardiol* 2000;86:610–4.
253. Engdahl J, Bang A, Lindqvist J, Herlitz J. Factors affecting shortand longterm prognosis among 1069 patients with outofhospital cardiac arrest and pulseless electrical activity. *Resuscitation* 2001;51:17–25.
254. Dumot JA, Burval DJ, Sprung J, et al. Outcome of adult cardiopulmonary resuscitations at a tertiary referral center including results of “limited” resuscitations. *Arch Intern Med* 2001;161:1751–8.
255. Tortolani AJ, Risucci DA, Powell SR, Dixon R. Inhospital cardiopulmonary resuscitation during asystole. Therapeutic factors associated with 24hour survival. *Chest* 1989;96:622–6.
256. Coon GA, Clinton JE, Ruiz E. Use of atropine for bradycardic prehospital cardiac arrest. *Ann Emerg Med* 1981;10:462–7.
257. Bottiger BW, Arntz HR, Chamberlain DA, et al. Thrombolysis during resuscitation for outofhospital cardiac arrest. *N Engl J Med* 2008;359:2651–62.
258. Böttiger BW, Martin E. Thrombolytic therapy during cardiopulmonary resuscitation and the role of coagulation activation after cardiac arrest. *Curr Opin Crit Care* 2001;7:176–83.
259. Spöhr F, Böttiger BW. Safety of thrombolysis during cardiopulmonary resuscitation. *Drug Saf* 2003;26:367–79.
260. Soar J, Foster J, Breikreutz R. Fluid infusion during CPR and after ROSC—is it safe? *Resuscitation* 2009;80:1221–2.
261. Price S, Uddin S, Quinn T. Echocardiography in cardiac arrest. *Curr Opin Crit Care* 2010;16:211–5.
262. Memtsoudis SG, Rosenberger P, Löffler M, et al. The usefulness of transesophageal echocardiography during intraoperative cardiac arrest in noncardiac surgery. *Anesth Analg* 2006;102:1653–7.
263. Comess KA, DeRook FA, Russell ML, TognazziEvans TA, Beach KW. The incidence of pulmonary embolism in unexplained sudden cardiac arrest with pulseless electrical activity. *Am J Med* 2000;109:351–6.
264. Niendorff DF, Rassias AJ, Palac R, Beach ML, Costa S, Greenberg M. Rapid cardiac ultrasound of inpatients suffering PEA arrest performed by nonexpert sonographers. *Resuscitation* 2005;67:81–7.
265. Tayal VS, Kline JA. Emergency echocardiography to detect pericardial effusion in patients in PEA and nearPEA states. *Resuscitation* 2003;59:315–8.
266. van der Wouw PA, Koster RW, Delemarre BJ, de Vos R, LampeSchoenmaeckers AJ, Lie KI. Diagnostic accuracy of transesophageal echocardiography during cardiopulmonary resuscitation. *J Am Coll Cardiol* 1997;30:780–3.
267. Hernandez C, Shuler K, Hannan H, Sonyika C, Likourezos A, Marshall J. C.A.U.S.E.: cardiac arrest ultrasound exam—a better approach to managing patients in primary nonarrhythmic cardiac arrest. *Resuscitation* 2008;76: 198–206.
268. Steiger HV, Rimbach K, Muller E, Breikreutz R. Focused emergency echocardiography: lifesaving tool for a 14yearold girl suffering outofhospital pulseless electrical activity arrest because of cardiac tamponade. *Eur J Emerg Med* 2009;16:103–5.
269. Breikreutz R, Walcher F, Seeger FH. Focused echocardiographic evaluation in resuscitation management: concept of an advanced life supportconformed algorithm. *Crit Care Med* 2007;35:S150–61.
270. Blaivas M, Fox JC. Outcome in cardiac arrest patients found to have cardiac standstill on the bedside emergency department echocardiogram. *Acad Emerg Med* 2001;8:616–21.
271. Salen P, O'Connor R, Sierzenski P, et al. Can cardiac sonography and capnography be used independently and in combination to predict resuscitation outcomes? *Acad Emerg Med* 2001;8:610–5.
272. Salen P, Melniker L, Chooljian C, et al. Does the presence or absence of sonographically identified cardiac activity predict resuscitation outcomes of cardiac arrest patients? *Am J Emerg Med* 2005;23:459–62.

273. Balan IS, Fiskum G, Hazelton J, CottoCumba C, Rosenthal RE. Oximetryguided reoxygenation improves neurological outcome after experimental cardiac arrest. *Stroke* 2006;37:3008–13.
274. Kilgannon JH, Jones AE, Shapiro NI, et al. Association between arterial hyperoxia following resuscitation from cardiac arrest and inhospital mortality. *JAMA* 2010;303:2165–71.
275. Nolan JP, Soar J. Airway techniques and ventilation strategies. *Curr Opin Crit Care* 2008;14:279–86.
276. GrmecS. Comparison of three different methods toconfirmtracheal tube placement in emergency intubation. *Intensive Care Med* 2002;28:701–4.
277. Lyon RM, Ferris JD, Young DM, McKeown DW, Oglesby AJ, Robertson C. Field intubation of cardiac arrest patients: a dying art? *Emerg Med J* 2010;27: 321–3.
- J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 1269
278. Jones JH, Murphy MP, Dickson RL, Somerville GG, Brizendine EJ. Emergency physicianverified outofhospital intubation: miss rates by paramedics. *Acad Emerg Med* 2004;11:707–9.
279. Pelucio M, Halligan L, Dhindsa H. Outofhospital experience with the syringe esophageal detector device. *Acad Emerg Med* 1997;4:563–8.
280. Jemmett ME, Kendal KM, Fourre MW, Burton JH. Unrecognized misplacement of endotracheal tubes in a mixed urban to rural emergency medical services setting. *Acad Emerg Med* 2003;10:961–5.
281. Katz SH, Falk JL. Misplaced endotracheal tubes by paramedics in an urban emergency medical services system. *Ann Emerg Med* 2001;37:32–7.
282. Wang HE, Simeone SJ, Weaver MD, Callaway CW. Interruptions in cardiopulmonary resuscitation from paramedic endotracheal intubation. *Ann Emerg Med* 2009;54:645–52, e1.
283. Gatward JJ, Thomas MJ, Nolan JP, Cook TM. Effect of chest compressions on the time taken to insert airway devices in a manikin. *Br J Anaesth* 2008;100: 351–6.
284. Li J. Capnography alone is imperfect for endotracheal tube placement confirmation during emergency intubation. *J Emerg Med* 2001;20:223–9.
285. Delguercio LR, Feins NR, Cohn JD, Coomaraswamy RP, Wollman SB, State D. Comparison of bloodflowduring external and internal cardiac massage in man. *Circulation* 1965;31(Suppl. 1):171–80.
286. Wik L, KramerJohansen J, Myklebust H, et al. Quality of cardiopulmonary resuscitation during outofhospital cardiac arrest. *JAMA* 2005;293:299–304.
287. KramerJohansen J, Myklebust H, Wik L, et al. Quality of outofhospital cardiopulmonary resuscitation with real time automated feedback: a prospective interventional study. *Resuscitation* 2006;71:283–92.
288. Sutton RM, Maltese MR, Niles D, et al. Quantitative analysis of chest compression interruptions during inhospital resuscitation of older children and adolescents. *Resuscitation* 2009;80:1259–63.
289. Sutton RM, Niles D, Nysaether J, et al. Quantitative analysis of CPR quality during inhospital resuscitation of older children and adolescents. *Pediatrics* 2009;124:494–9.
290. Cabrini L, Beccaria P, Landoni G, et al. Impact of impedance threshold devices on cardiopulmonary resuscitation: a systematic review and metaanalysis of randomized controlled studies. *Crit Care Med* 2008;36:1625–32.
291. Steen S, Liao Q, Pierre L, Paskevicius A, Sjoberg T. Evaluation of LUCAS, a new device for automatic mechanical compression and active decompression resuscitation. *Resuscitation* 2002;55:285–99.
292. Rubertsson S, Karlsten R. Increased cortical cerebral blood flow with LUCAS; a new device for mechanical chest compressions compared to standard external compressions during experimental cardiopulmonary resuscitation. *Resuscitation* 2005;65:357–63.
293. Timerman S, Cardoso LF, Ramires JA, Halperin H. Improved hemodynamic performance with a novel chest compression device during treatment of inhospital cardiac arrest. *Resuscitation* 2004;61:273–80.
294. Halperin H, Berger R, Chandra N, et al. Cardiopulmonary resuscitation with a hydraulicpneumatic band. *Crit Care Med* 2000;28:N203–6.
295. Halperin HR, Paradis N, Ornato JP, et al. Cardiopulmonary resuscitation with a novel chest compression device in a porcine model of cardiac arrest: improved hemodynamics and mechanisms. *J Am Coll Cardiol* 2004;44:2214–20.
296. Hallstrom A, Rea TD, Sayre MR, et al. Manual chest compression vs use of an automated chest compression device during resuscitation following outofhospital cardiac arrest: a randomized trial. *JAMA* 2006;295:2620–8.
297. Ong ME, Ornato JP, Edwards DP, et al. Use of an automated, loadistributing band chest compression device for outofhospital cardiac arrest resuscitation. *JAMA* 2006;295:2629–37.
298. Larsen AI, Hjernevik AS, Ellingsen CL, Nilsen DW. Cardiac arrest with continuous mechanical chest compression during percutaneous coronary intervention. A report on the use of the LUCAS device. *Resuscitation* 2007;75:454–9.

299. Wagner H, Terkelsen CJ, Friberg H, et al. Cardiac arrest in the catheterisation laboratory: a 5 year experience of using mechanical chest compressions to facilitate PCI during prolonged resuscitation efforts. *Resuscitation* 2010;81:383–7.
300. Wirth S, Korner M, Treitl M, et al. Computed tomography during cardiopulmonary resuscitation using automated chest compression devices—an initial study. *Eur Radiol* 2009;19:1857–66.
301. Holmstrom P, Boyd J, Sorsa M, Kuisma M. A case of hypothermic cardiac arrest treated with an external chest compression device (LUCAS) during transport to rewarming. *Resuscitation* 2005;67:139–41.
302. Wik L, Kiil S. Use of an automatic mechanical chest compression device (LUCAS) as a bridge to establishing cardiopulmonary bypass for a patient with hypothermic cardiac arrest. *Resuscitation* 2005;66:391–4.
303. Olasveengen TM, Wik L, Steen PA. Quality of cardiopulmonary resuscitation before and during transport in outofhospital cardiac arrest. *Resuscitation* 2008;76:185–90.
304. Sunde K, Wik L, Steen PA. Quality of mechanical, manual standard and active compression–decompression CPR on the arrest site and during transport in a manikin model. *Resuscitation* 1997;34:235–42.
305. Laver S, Farrow C, Turner D, Nolan J. Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive Care Med* 2004;30:2126–8.
306. Laurent I, Monchi M, Chiche JD, et al. Reversible myocardial dysfunction in survivors of outofhospital cardiac arrest. *J Am Coll Cardiol* 2002;40:2110–6.
307. RuizBailen M, Aguayo de Hoyos E, RuizNavarro S, et al. Reversible myocardial dysfunction after cardiopulmonary resuscitation. *Resuscitation* 2005;66:175–81.
308. Cerchiari EL, Safar P, Klein E, Diven W. Visceral, hematologic and bacteriologic changes and neurologic outcome after cardiac arrest in dogs. The visceral postresuscitation syndrome. *Resuscitation* 1993;25:119–36.
309. Adrie C, Monchi M, Laurent I, et al. Coagulopathy after successful cardiopulmonary resuscitation following cardiac arrest: implication of the protein C anticoagulant pathway. *J Am Coll Cardiol* 2005;46:21–8.
310. Adrie C, AdibConquy M, Laurent I, et al. Successful cardiopulmonary resuscitation after cardiac arrest as a “sepsislike” syndrome. *Circulation* 2002;106:562–8.
311. Adrie C, Laurent I, Monchi M, Cariou A, Dhainaut JF, Spaulding C. Postresuscitation disease after cardiac arrest: a sepsislike syndrome? *Curr Opin Crit Care* 2004;10:208–12.
312. Zwemer CF, Whitesall SE, D’Alecry LG. Cardiopulmonary cerebral resuscitation with 100% oxygen exacerbates neurological dysfunction following nine minutes of normothermic cardiac arrest in dogs. *Resuscitation* 1994;27:159–70.
313. Richards EM, Fiskum G, Rosenthal RE, Hopkins I, McKenna MC. Hyperoxic reperfusion after global ischemia decreases hippocampal energy metabolism. *Stroke* 2007;38:1578–84.
314. Vereczki V, Martin E, Rosenthal RE, Hof PR, Hoffman GE, Fiskum G. Normoxic resuscitation after cardiac arrest protects against hippocampal oxidative stress, metabolic dysfunction, and neuronal death. *J Cereb Blood Flow Metab* 2006;26:821–35.
315. Liu Y, Rosenthal RE, Haywood Y, MiljkovicLolic M, Vanderhoek JY, Fiskum G. Normoxic ventilation after cardiac arrest reduces oxidation of brain lipids and improves neurological outcome. *Stroke* 1998;29:1679–86.
316. Spaulding CM, Joly LM, Rosenberg A, et al. Immediate coronary angiography in survivors of outofhospital cardiac arrest. *N Engl J Med* 1997;336:1629–33.
317. Sunde K, Pytte M, Jacobsen D, et al. Implementation of a standardised treatment protocol for post resuscitation care after outofhospital cardiac arrest. *Resuscitation* 2007;73:29–39.
318. Bendz B, Eritsland J, Nakstad AR, et al. Longterm prognosis after outofhospital cardiac arrest and primary percutaneous coronary intervention. *Resuscitation* 2004;63:49–53.
319. Keelan PC, Bunch TJ, White RD, Packer DL, Holmes Jr DR. Early direct coronary angioplasty in survivors of outofhospital cardiac arrest. *Am J Cardiol* 2003;91:1461–3. A6.
320. QuinteroMoran B, Moreno R, Villarreal S, et al. Percutaneous coronary intervention for cardiac arrest secondary to STelevation acute myocardial infarction. Influence of immediate paramedical/medical assistance on clinical outcome. *J Invasive Cardiol* 2006;18:269–72.
321. Garot P, Lefevre T, Eltchaninoff H, et al. Sixmonth outcome of emergency percutaneous coronary intervention in resuscitated patients after cardiac arrest complicating STelevation myocardial infarction. *Circulation* 2007;115:1354–62.
322. Nagao K, Hayashi N, Kanmatsuse K, et al. Cardiopulmonary cerebral resuscitation using emergency cardiopulmonary bypass, coronary reperfusion therapy and mild hypothermia in patients with cardiac arrest outside the hospital. *J Am Coll Cardiol* 2000;36:776–83.

323. Knafelj R, Radsel P, Ploj T, Noc M. Primary percutaneous coronary intervention and mild induced hypothermia in comatose survivors of ventricular fibrillation with STelevation acute myocardial infarction. *Resuscitation* 2007;74: 227–34.
324. Nielsen N, Hovdenes J, Nilsson F, et al. Outcome, timing and adverse events in therapeutic hypothermia after outofhospital cardiac arrest. *Acta Anaesthesiol Scand* 2009;53:926–34.
325. Hovdenes J, Laake JH, Aaberge L, Haugaa H, Bugge JF. Therapeutic hypothermia after outofhospital cardiac arrest: experiences with patients treated with percutaneous coronary intervention and cardiogenic shock. *Acta Anaesthesiol Scand* 2007;51:137–42.
326. Wolfrum S, Pierau C, Radke PW, Schunkert H, Kurowski V. Mild therapeutic hypothermia in patients after outofhospital cardiac arrest due to acute STsegment elevation myocardial infarction undergoing immediate percutaneous coronary intervention. *Crit Care Med* 2008;36:1780–6.
327. Snyder BD, Hauser WA, Loewenson RB, Leppik IE, RamirezLassepas M, Gumnit RJ. Neurologic prognosis after cardiopulmonary arrest. III. Seizure activity. *Neurology* 1980;30:1292–7.
328. Levy DE, Caronna JJ, Singer BH, Lapinski RH, Frydman H, Plum F. Predicting outcome from hypoxicischemic coma. *JAMA* 1985;253:1420–6.
329. Krumholz A, Stern BJ, Weiss HD. Outcome from coma after cardiopulmonary resuscitation: relation to seizures and myoclonus. *Neurology* 1988;38:401–5.
330. Zandbergen EG, Hijdra A, Koelman JH, et al. Prediction of poor outcome within the first 3 days of postanoxic coma. *Neurology* 2006;66:62–8.
331. Ingvar M. Cerebral blood flow and metabolic rate during seizures. Relationship to epileptic brain damage. *Ann N Y Acad Sci* 1986;462:194–206.
332. Nolan JP, Laver SR, Welch CA, Harrison DA, Gupta V, Rowan K. Outcome following admission to UK intensive care units after cardiac arrest: a secondary analysis of the ICNARC Case Mix Programme Database. *Anaesthesia* 2007;62:1207–16.
333. Losert H, Sterz F, Roine RO, et al. Strict normoglycaemic blood glucose levels in the therapeutic management of patients within 12 h after cardiac arrest might not be necessary. *Resuscitation* 2007.
334. Skrifvars MB, Saarinen K, Ikola K, Kuisma M. Improved survival after inhospital cardiac arrest outside critical care areas. *Acta Anaesthesiol Scand* 2005;49:1534–9.
335. Mullner M, Sterz F, Binder M, Schreiber W, Deimel A, Lagner AN. Blood glucose concentration after cardiopulmonary resuscitation influences functional 1270 *J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* neurological recovery in human cardiac arrest survivors. *J Cereb Blood Flow Metab* 1997;17:430–6.
336. Calle PA, Buylaert WA, Vanhaute OA. Glycemia in the postresuscitation period. The Cerebral Resuscitation Study Group. *Resuscitation* 1989;17(Suppl.):S181–8 [discussion S99–S206].
337. Longstreth Jr WT, Diehr P, Inui TS. Prediction of awakening after outofhospital cardiac arrest. *N Engl J Med* 1983;308:1378–82.
338. Longstreth Jr WT, Inui TS. High blood glucose level on hospital admission and poor neurological recovery after cardiac arrest. *Ann Neurol* 1984;15:59–63.
339. Finfer S, Chittock DR, Su SY, et al. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;360:1283–97.
340. Preiser JC, Devos P, RuizSantana S, et al. A prospective randomised multicentre controlled trial on tight glucose control by intensive insulin therapy in adult intensive care units: the Glucontrol study. *Intensive Care Med* 2009;35:1738–48.
341. Griesdale DE, de Souza RJ, van Dam RM, et al. Intensive insulin therapy and mortality among critically ill patients: a metaanalysis including NICESUGAR study data. *CMAJ* 2009;180:821–7.
342. Wiener RS, Wiener DC, Larson RJ. Benefits and risks of tight glucose control in critically ill adults: a metaanalysis. *JAMA* 2008;300:933–44.
343. Krinsley JS, Grover A. Severe hypoglycemia in critically ill patients: risk factors and outcomes. *Crit Care Med* 2007;35:2262–7.
344. Meyfroidt G, Keenan DM, Wang X, Wouters PJ, Veldhuis JD, Van den Berghe G. Dynamic characteristics of blood glucose time series during the course of critical illness: effects of intensive insulin therapy and relative association with mortality. *Crit Care Med* 2010;38:1021–9.
345. Padkin A. Glucose control after cardiac arrest. *Resuscitation* 2009;80:611–2.
346. Takino M, Okada Y. Hyperthermia following cardiopulmonary resuscitation. *Intensive Care Med* 1991;17:419–20.
347. Hickey RW, Kochanek PM, Ferimer H, Alexander HL, Garman RH, Graham SH. Induced hyperthermia exacerbates neurologic neuronal histologic damage after asphyxial cardiac arrest in rats. *Crit Care Med* 2003;31:531–5.

348. Takasu A, Saitoh D, Kaneko N, Sakamoto T, Okada Y. Hyperthermia: is it an ominous sign after cardiac arrest? *Resuscitation* 2001;49:273–7.
349. Zeiner A, Holzer M, Sterz F, et al. Hyperthermia after cardiac arrest is associated with an unfavorable neurologic outcome. *Arch Intern Med* 2001;161:2007–12.
350. Hickey RW, Kochanek PM, Ferimer H, Graham SH, Safar P. Hypothermia and hyperthermia in children after resuscitation from cardiac arrest. *Pediatrics* 2000;106:118–22.
351. Diringer MN, Reaven NL, Funk SE, Uman GC. Elevated body temperature independently contributes to increased length of stay in neurologic intensive care unit patients. *Crit Care Med* 2004;32:1489–95.
352. Gunn AJ, Thoresen M. Hypothermic neuroprotection. *NeuroRx* 2006;3:154–69.
353. Froehler MT, Geocadin RG. Hypothermia for neuroprotection after cardiac arrest: mechanisms, clinical trials and patient care. *J Neurol Sci* 2007;261:118–26.
354. McCullough JN, Zhang N, Reich DL, et al. Cerebral metabolic suppression during hypothermic circulatory arrest in humans. *Ann Thorac Surg* 1999;67:1895–9 [discussion 919–21].
355. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;346:549–56.
356. Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of outofhospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;346:557–63.
357. Bernard SA, Jones BM, Horne MK. Clinical trial of induced hypothermia in comatose survivors of outofhospital cardiac arrest. *Ann Emerg Med* 1997;30:146–53.
358. Oddo M, Schaller MD, Feihl F, Ribordy V, Liaudet L. From evidence to clinical practice: effective implementation of therapeutic hypothermia to improve patient outcome after cardiac arrest. *Crit Care Med* 2006;34:1865–73.
359. Busch M, Soreide E, Lossius HM, Lexow K, Dickstein K. Rapid implementation of therapeutic hypothermia in comatose outofhospital cardiac arrest survivors. *Acta Anaesthesiol Scand* 2006;50:1277–83.
360. Storm C, Steffen I, Schefold JC, et al. Mild therapeutic hypothermia shortens intensive care unit stay of survivors after outofhospital cardiac arrest compared to historical controls. *Crit Care* 2008;12:R78.
361. Don CW, Longstreth Jr WT, Maynard C, et al. Active surface cooling protocol to induce mild therapeutic hypothermia after outofhospital cardiac arrest: a retrospective beforeandafter comparison in a single hospital. *Crit Care Med* 2009;37:3062–9.
362. Arrich J. Clinical application of mild therapeutic hypothermia after cardiac arrest. *Crit Care Med* 2007;35:1041–7.
363. Holzer M, Mullner M, Sterz F, et al. Efficacy and safety of endovascular cooling after cardiac arrest: cohort study and Bayesian approach. *Stroke* 2006;37:1792–7.
364. Polderman KH, Herold I. Therapeutic hypothermia and controlled normothermia in the intensive care unit: practical considerations, side effects, and cooling methods. *Crit Care Med* 2009;37:1101–20.
365. Kuboyama K, Safar P, Radovsky A, et al. Delay in cooling negates the beneficial effect of mild resuscitative cerebral hypothermia after cardiac arrest in dogs: a prospective, randomized study. *Crit Care Med* 1993;21:1348–58.
366. Edgren E, Hedstrand U, Nordin M, Rydin E, Ronquist G. Prediction of outcome after cardiac arrest. *Crit Care Med* 1987;15:820–5.
367. Young GB, Doig G, Ragazzoni A. Anoxicischemic encephalopathy: clinical and electrophysiological associations with outcome. *Neurocrit Care* 2005;2:159–64.
368. Al Thenayan E, Savard M, Sharpe M, Norton L, Young B. Predictors of poor neurologic outcome after induced mild hypothermia following cardiac arrest. *Neurology* 2008;71:1535–7.
369. Wijdicks EF, Parisi JE, Sharbrough FW. Prognostic value of myoclonus status in comatose survivors of cardiac arrest. *Ann Neurol* 1994;35:239–43.
370. Thomke F, Marx JJ, Sauer O, et al. Observations on comatose survivors of cardiopulmonary resuscitation with generalized myoclonus. *BMC Neurol* 2005;5:14.
371. Arnoldus EP, Lammers GJ. Postanoxic coma: good recovery despite myoclonus status. *Ann Neurol* 1995;38:697–8.
372. Celesia GG, Grigg MM, Ross E. Generalized status myoclonicus in acute anoxic and toxicmetabolic encephalopathies. *Arch Neurol* 1988;45:781–4.
373. Morris HR, Howard RS, Brown P. Early myoclonic status and outcome after cardiorespiratory arrest. *J Neurol Neurosurg Psychiatry* 1998;64: 267–8.
374. Datta S, Hart GK, Opdam H, Gutteridge G, Archer J. Posthypoxic myoclonic status: the prognosis is not always hopeless. *Crit Care Resusc* 2009;11:39–41.
375. English WA, Giffin NJ, Nolan JP. Myoclonus after cardiac arrest: pitfalls in diagnosis and prognosis. *Anaesthesia* 2009;64:908–11.

376. Wijdicks EF, Hijdra A, Young GB, Bassetti CL, Wiebe S. Practice parameter: prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidencebased review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006;67:203–10.
377. Tiainen M, Kovala TT, Takkunen OS, Roine RO. Somatosensory and brainstem auditory evoked potentials in cardiac arrest patients treated with hypothermia. *Crit Care Med* 2005;33:1736–40.
378. Rossetti AO, Oddo M, Liaudet L, Kaplan PW. Predictors of awakening from postanoxic status epilepticus after therapeutic hypothermia. *Neurology* 2009;72:744–9.
379. Rossetti AO, Logroscino G, Liaudet L, et al. Status epilepticus: an independent outcome predictor after cerebral anoxia. *Neurology* 2007;69:255–60.
380. Fioux F, Losser MR, Bourgeois E, et al. Kidney retrieval after sudden out of hospital refractory cardiac arrest: a cohort of uncontrolled non heart beating donors. *Crit Care* 2009;13:R141.
381. Kootstra G. Statement on nonheartbeating donor programs. *Transplant Proc* 1995;27:2965.
382. Vermeer F, Oude Ophuis AJ, vd Berg EJ, et al. Prospective randomised comparison between thrombolysis, rescue PTCA, and primary PTCA in patients with extensive myocardial infarction admitted to a hospital without PTCA facilities: a safety and feasibility study. *Heart* 1999;82:426–31.
383. Widimsky P, Groch L, Zelizko M, Aschermann M, Bednar F, Suryapranata H. Multicentre randomized trial comparing transport to primary angioplasty vs immediate thrombolysis vs combined strategy for patients with acute myocardial infarction presenting to a community hospital without a catheterization laboratory. The PRAGUE study. *Eur Heart J* 2000;21:823–31.
384. Widimsky P, Budesinsky T, Vorac D, et al. Long distance transport for primary angioplasty vs immediate thrombolysis in acute myocardial infarction. Final results of the randomized national multicentre trial—PRAGUE2. *Eur Heart J* 2003;24:94–104.
385. Le May MR, So DY, Dionne R, et al. A citywide protocol for primary PCI in STsegment elevation myocardial infarction. *N Engl J Med* 2008;358:231–40.
386. Abernathy 3rd JH, McGwin Jr G, Acker 3rd JE, Rue 3rd LW. Impact of a voluntary trauma system on mortality, length of stay, and cost at a level I trauma center. *Am Surg* 2002;68:182–92.
387. Clemmer TP, Orme Jr JF, Thomas FO, Brooks KA. Outcome of critically injured patients treated at Level I trauma centers versus fullservice community hospitals. *Crit Care Med* 1985;13:861–3.
388. Culica D, Aday LA, Rohrer JE. Regionalized trauma care system in Texas: implications for redesigning trauma systems. *Med Sci Monit* 2007;13:SR9–18.
389. Hannan EL, Farrell LS, Cooper A, Henry M, Simon B, Simon R. Physiologic trauma triage criteria in adult trauma patients: are they effective in saving lives by transporting patients to trauma centers? *J Am Coll Surg* 2005;200:584–92.
390. Harrington DT, Connolly M, Biffi WL, Majercik SD, Cioffi WG. Transfer times to definitive care facilities are too long: a consequence of an immature trauma system. *Ann Surg* 2005;241:961–6 [discussion 6–8].
391. Liberman M, Mulder DS, Lavoie A, Sampalis JS. Implementation of a trauma care system: evolution through evaluation. *J Trauma* 2004;56:1330–5.
392. MacKenzie EJ, Rivara FP, Jurkovich GJ, et al. A national evaluation of the effect of traumacenter care on mortality. *N Engl J Med* 2006;354:366–78.
393. Mann NC, Cahn RM, Mullins RJ, Brand DM, Jurkovich GJ. Survival among injured geriatric patients during construction of a statewide trauma system. *J Trauma* 2001;50:1111–6.
394. Mullins RJ, VeumStone J, Hedges JR, et al. Influence of a statewide trauma system on location of hospitalization and outcome of injured patients. *J Trauma* 1996;40:536–45 [discussion 45–46].
395. Mullins RJ, Mann NC, Hedges JR, Worrall W, Jurkovich GJ. Preferential benefit of implementation of a statewide trauma system in one of two adjacent states. *J Trauma* 1998;44:609–16 [discussion 17].
396. Mullins RJ, VeumStone J, Helfand M, et al. Outcome of hospitalized injured patients after institution of a trauma system in an urban area. *JAMA* 1994;271:1919–24.
397. Mullner R, Goldberg J. An evaluation of the Illinois trauma system. *Med Care* 1978;16:140–51.
398. Mullner R, Goldberg J. Toward an outcomeoriented medical geography: an evaluation of the Illinois trauma/emergency medical services system. *Soc Sci Med* 1978;12:103–10.
- J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 1271
399. Nathens AB, Jurkovich GJ, Rivara FP, Maier RV. Effectiveness of state trauma systems in reducing injuryrelated mortality: a national evaluation. *J Trauma* 2000;48:25–30 [discussion 1].
400. Nathens AB, Maier RV, Brundage SI, Jurkovich GJ, Grossman DC. The effect of interfacility transfer on outcome in an urban trauma system. *J Trauma* 2003;55:444–9.
401. Nicholl J, Turner J. Effectiveness of a regional trauma system in reducing mortality from major trauma: before and after study. *BMJ* 1997;315:1349–54.

402. Potoka DA, Schall LC, Gardner MJ, Stafford PW, Peitzman AB, Ford HR. Impact of pediatric trauma centers on mortality in a statewide system. *J Trauma* 2000;49:237–45.
403. Sampalis JS, Lavoie A, Boukas S, et al. Trauma center designation: initial impact on traumarelated mortality. *J Trauma* 1995;39:232–7 [discussion 7–9].
404. Sampalis JS, Denis R, Frechette P, Brown R, Fleiszer D, Mulder D. Direct transport to tertiary trauma centers versus transfer from lower level facilities: impact on mortality and morbidity among patients with major trauma. *J Trauma* 1997;43:288–95 [discussion 95–96].
405. Nichol G, Aufderheide TP, Eigel B, et al. Regional systems of care for outofhospital cardiac arrest: a policy statement from the American Heart Association. *Circulation* 2010;121:709–29.
406. Nichol G, Soar J. Regional cardiac resuscitation systems of care. *Curr Opin Crit Care* 2010;16:223–30.
407. Soar J, Packham S. Cardiac arrest centres make sense. *Resuscitation* 2010;81:507–8.
408. TunstallPedoe H, Vanuzzo D, Hobbs M, et al. Estimation of contribution of changes in coronary care to improving survival, event rates, and coronary heart disease mortality across the WHO MONICA Project populations. *Lancet* 2000;355:688–700.
409. Fox KA, Cokkinos DV, Deckers J, Keil U, Maggioni A, Steg G. The ENACT study: a panEuropean survey of acute coronary syndromes. European Network for Acute Coronary Treatment. *Eur Heart J* 2000;21:1440–9.
410. Goodman SG, Huang W, Yan AT, et al. The expanded global registry of acute coronary events: baseline characteristics, management practices, and hospital outcomes of patients with acute coronary syndromes. *Am Heart J* 2009;158:193–201, e1–e5.
411. Lowel H, Meisinger C, Heier M, et al. Sex specific trends of sudden cardiac death and acute myocardial infarction: results of the populationbased KORA/MONICA Augsburg register 1985 to 1998. *Dtsch Med Wochenschr* 2002;127:2311–6.
412. Thygesen K, Alpert JS, White HD. Universal definition of myocardial infarction. *Eur Heart J* 2007;28:2525–38.
413. Van de Werf F, Bax J, Betriu A, et al. Management of acute myocardial infarction in patients presenting with persistent STsegment elevation: the Task Force on the Management of STSegment Elevation Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J* 2008;29:2909–45.
414. Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with STElevation myocardial infarction—executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 1999 Guidelines for the Management of Patients With Acute Myocardial Infarction). *Circulation* 2004;110:588–636.
415. Douglas PS, Ginsburg GS. The evaluation of chest pain in women. *N Engl J Med* 1996;334:1311–5.
416. Solomon CG, Lee TH, Cook EF, et al. Comparison of clinical presentation of acute myocardial infarction in patients older than 65 years of age to younger patients: the Multicenter Chest Pain Study experience. *Am J Cardiol* 1989;63: 772–6.
417. Ioannidis JP, Salem D, Chew PW, Lau J. Accuracy and clinical effect of outofhospital electrocardiography in the diagnosis of acute cardiac ischemia: a metaanalysis. *Ann Emerg Med* 2001;37:461–70.
418. Kudenchuk PJ, Ho MT, Weaver WD, et al. Accuracy of computerinterpreted electrocardiography in selecting patients for thrombolytic therapy. MITI Project Investigators. *J Am Coll Cardiol* 1991;17:1486–91.
419. Dhruva VN, Abdelhadi SI, Anis A, et al. STSegment Analysis Using Wireless Technology in Acute Myocardial Infarction (STATMI) trial. *J Am Coll Cardiol* 2007;50:509–13.
420. AntmanEM, Tanasijevic MJ,ThompsonB, et al. Cardiacspecific troponin I levels to predict the risk of mortality in patients with acute coronary syndromes. *N Engl J Med* 1996;335:1342–9.
421. Hess EP, Thiruganasambandamoorthy V, Wells GA, et al. Diagnostic accuracy of clinical prediction rules to exclude acute coronary syndrome in the emergency department setting: a systematic review. *CJEM* 2008;10:373–82.
422. Ramakrishna G, Milavetz JJ, Zinsmeister AR, et al. Effect of exercise treadmill testing and stress imaging on the triage of patients with chest pain: CHEER substudy. *Mayo Clin Proc* 2005;80:322–9.
423. Kearney PM, Baigent C, Godwin J, Halls H, Emberson JR, Patrono C. Do selective cyclooxygenase2 inhibitors and traditional nonsteroidal antiinflammatory drugs increase the risk of atherothrombosis? Metaanalysis of randomised trials. *BMJ* 2006;332:1302–8.
424. Rawles JM, Kenmure AC. Controlled trial of oxygen in uncomplicated myocardial infarction. *BMJ* 1976;1:1121–3.
425. Wijesinghe M, Perrin K, Ranchord A, Simmonds M, Weatherall M, Beasley R. Routine use of oxygen in the treatment of myocardial infarction: systematic review. *Heart* 2009;95:198–202.

426. Cabello JB, Burls A, Emparanza JI, Bayliss S, Quinn T. Oxygen therapy for acute myocardial infarction. *Cochrane Database Syst Rev* 2010;6:CD007160.
427. O'Driscoll BR, Howard LS, Davison AG. BTS guideline for emergency oxygen use in adult patients. *Thorax* 2008;63(Suppl. 6):vi1-68.
428. Freimark D, Matetzky S, Leor J, et al. Timing of aspirin administration as a determinant of survival of patients with acute myocardial infarction treated with thrombolysis. *Am J Cardiol* 2002;89:381-5.
429. Barbash IM, Freimark D, Gottlieb S, et al. Outcome of myocardial infarction in patients treated with aspirin is enhanced by prehospital administration *Cardiology* 2002;98:141-7.
430. Yusuf S, Zhao F, Mehta SR, Chrolavicius S, Tognoni G, Fox KK. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without STsegment elevation. *N Engl J Med* 2001;345:494-502.
431. Mehta SR, Yusuf S, Peters RJ, et al. Effects of pretreatment with clopidogrel and aspirin followed by longterm therapy in patients undergoing percutaneous coronary intervention: the PCICURE study. *Lancet* 2001;358: 527-33.
432. TIMI11B Investigators, Antman EM, McCabe CH, et al. Enoxaparin prevents death and cardiac ischemic events in unstable angina/nonQwave myocardial infarction. Results of the thrombolysis in myocardial infarction (TIMI) 11B trial. *Circulation* 1999;100:1593-601.
433. Cohen M, Demers C, Gurfinkel EP, et al. A comparison of lowmolecularweight heparin with unfractionated heparin for unstable coronary artery disease. Efficacy and safety of subcutaneous enoxaparin in nonQwave coronary events study group. *N Engl J Med* 1997;337:447-52.
434. Yusuf S, Mehta SR, Chrolavicius S, et al. Comparison of fondaparinux and enoxaparin in acute coronary syndromes. *N Engl J Med* 2006;354:1464-76.
435. Mehta SR, Boden WE, Eikelboom JW, et al. Antithrombotic therapy with fondaparinux in relation to interventional management strategy in patients with STand nonSTsegment elevation acute coronary syndromes: an individual patientlevel combined analysis of the Fifth and Sixth Organization to Assess Strategies in Ischemic Syndromes (OASIS 5 and 6) randomized trials. *Circulation* 2008;118:2038-46.
436. Lincoff AM, Bittl JA, Harrington RA, et al. Bivalirudin and provisional glycoprotein IIb/IIIa blockade compared with heparin and planned glycoprotein IIb/IIIa blockade during percutaneous coronary intervention: REPLACE2 randomized trial. *JAMA* 2003;289:853-63.
437. Efficacy and safety of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin: the ASSENT3 randomised trial in acute myocardial infarction. *Lancet* 2001;358:605-13.
438. Eikelboom JW, Quinlan DJ, Mehta SR, Turpie AG, Menown IB, Yusuf S. Unfractionated and lowmolecularweight heparin as adjuncts to thrombolysis in aspirintreated patients with STElevation acute myocardial infarction: a metaanalysis of the randomized trials. *Circulation* 2005;112:3855-67.
439. Wallentin L, Goldstein P, Armstrong PW, et al. Efficacy and safety of tenecteplase in combination with the lowmolecularweight heparin enoxaparin or unfractionated heparin in the prehospital setting: the Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)3 PLUS randomized trial in acute myocardial infarction. *Circulation* 2003;108: 135-42.
440. Zeymer U, Gitt A, Junger C, et al. Efficacy and safety of enoxaparin in unselected patients with STsegment elevation myocardial infarction. *Thromb Haemost* 2008;99:150-4.
441. Zeymer U, Gitt A, Zahn R, et al. Efficacy and safety of enoxaparin in combination with and without GP IIb/IIIa inhibitors in unselected patients with ST segment elevation myocardial infarction treated with primary percutaneous coronary intervention. *EuroIntervention* 2009;4:524-8.
442. Bassand JP, Hamm CW, Ardissino D, et al. Guidelines for the diagnosis and treatment of nonSTsegment elevation acute coronary syndromes. *Eur Heart J* 2007;28:1598-660.
443. Anderson JL, Adams CD, Antman EM, et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non STElevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non STElevation Myocardial Infarction): developed in collaboration with the American College of Emergency Physicians, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons: endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation and the Society for Academic Emergency Medicine. *Circulation* 2007;116:e148-304.
444. Kushner FG, Hand M, Smith SCJr, et al. 2009 Focused Updates: ACC/AHA guidelines for the management of patients with STElevation myocardial infarction (updating the 2004 Guideline and 2007 Focused Update) and ACC/AHA/SCAI Guidelines on Percutaneous Coronary Intervention (updating the 2005 Guideline and 2007 Focused Update): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2009;120:2271-306. Erratum in: *Circulation*. 010 March 30;121(12):e257. Dosage error in article text. 445. Boersma E, Maas AC,

Deckers JW, Simoons ML. Early thrombolytic treatment in acute myocardial infarction: reappraisal of the golden hour. *Lancet* 1996;348:771–5.

446. Prehospital thrombolytic therapy in patients with suspected acute myocardial infarction. The European Myocardial Infarction Project Group. *N Engl J Med* 1993;329:383–9.

447. Weaver WD, Cerqueira M, Hallstrom AP, et al. Prehospitalinitiated vs hospitalinitiated thrombolytic therapy. The Myocardial Infarction Triage and Intervention Trial. *JAMA* 1993;270:1211–6.

1272 J.P. Nolan et al. / *Resuscitation* 81 (2010) 1219–1276 448. Feasibility, safety, and efficacy of domiciliary thrombolysis by general practitioners: Grampian region early anistreplase trial. GREAT Group. *BMJ* 1992;305:548–53.

449. Welsh RC, Travers A, Senaratne M, Williams R, Armstrong PW. Feasibility and applicability of paramedicbased prehospital fibrinolysis in a large North American center. *Am Heart J* 2006;152:1007–14.

450. Pedley DK, Bissett K, Connolly EM, et al. Prospective observational cohort study of time saved by prehospital thrombolysis for ST elevation myocardial infarction delivered by paramedics. *BMJ* 2003;327:22–6.

451. Grijseels EW, Bouten MJ, Lenderink T, et al. Prehospital thrombolytic therapy with either alteplase or streptokinase. Practical applications, complications and longterm results in 529 patients. *Eur Heart J* 1995;16:1833–8.

452. Morrison LJ, Verbeek PR, McDonald AC, Sawadsky BV, Cook DJ. Mortality and prehospital thrombolysis for acute myocardial infarction: a metaanalysis. *JAMA* 2000;283:2686–92.

453. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet* 2003;361:13–20.

454. Dalby M, Bouzamondo A, Lechat P, Montalescot G. Transfer for primary angioplasty versus immediate thrombolysis in acute myocardial infarction: a metaanalysis. *Circulation* 2003;108:1809–14.

455. Steg PG, Bonnefoy E, Chabaud S, et al. Impact of time to treatment on mortality after prehospital fibrinolysis or primary angioplasty: data from the CAPTIM randomized clinical trial. *Circulation* 2003;108:2851–6.

456. Bonnefoy E, Steg PG, Boutitie F, et al. Comparison of primary angioplasty and prehospital fibrinolysis in acute myocardial infarction (CAPTIM) trial: a 5year followup. *Eur Heart J* 2009;30:1598–606.

457. Kalla K, Christ G, Karnik R, et al. Implementation of guidelines improves the standard of care: the Viennese registry on reperfusion strategies in STElevation myocardial infarction (Vienna STEMI registry). *Circulation* 2006;113:2398–405.

458. Primary versus tenecteplasefacilitated percutaneous coronary intervention in patients with STsegment elevation acute myocardial infarction (ASSENT4 PCI): randomised trial. *Lancet* 2006;367:569–78.

459. Pinto DS, Kirtane AJ, Nallamothu BK, et al. Hospital delays in reperfusion for STElevation myocardial infarction: implications when selecting a reperfusion strategy. *Circulation* 2006;114:2019–25.

460. Keeley EC, Boura JA, Grines CL. Comparison of primary and facilitated percutaneous coronary interventions for STElevation myocardial infarction: quantitative review of randomised trials. *Lancet* 2006;367:579–88.

461. Herrmann HC, Lu J, Brodie BR, et al. Benefit of facilitated percutaneous coronary intervention in highrisk STsegment elevation myocardial infarction patients presenting to nonpercutaneous coronary intervention hospitals. *JACC Cardiovasc Interv* 2009;2:917–24.

462. Gershlick AH, StephensLloyd A, Hughes S, et al. Rescue angioplasty after failed thrombolytic therapy for acute myocardial infarction. *N Engl J Med* 2005;353:2758–68.

463. Danchin N, Coste P, Ferrieres J, et al. Comparison of thrombolysis followed by broad use of percutaneous coronary intervention with primary percutaneous coronary intervention for STsegmentelevation acute myocardial infarction: data from the french registry on acute STElevation myocardial infarction (FASTMI). *Circulation* 2008;118:268–76.

464. Cantor WJ, Fitchett D, Borgundvaag B, et al. Routine early angioplasty after fibrinolysis for acute myocardial infarction. *N Engl J Med* 2009;360:2705–18.

465. Redding JS. The choking controversy: critique of evidence on the Heimlich maneuver. *Crit Care Med* 1979;7:475–9.

466. Kuisma M, Suominen P, Korpela R. Paediatric outofhospital cardiac arrests—epidemiology and outcome. *Resuscitation* 1995;30:141–50.

467. Sirbaugh PE, Pepe PE, Shook JE, et al. A prospective, populationbased study of the demographics, epidemiology, management, and outcome of outofhospital pediatric cardiopulmonary arrest. *Ann Emerg Med* 1999;33:174–84.

468. Hickey RW, Cohen DM, Strausbaugh S, Dietrich AM. Pediatric patients requiring CPR in the prehospital setting. *Ann Emerg Med* 1995;25:495–501.

469. Young KD, Seidel JS. Pediatric cardiopulmonary resuscitation: a collective review. *Ann Emerg Med* 1999;33:195–205.
470. Reis AG, Nadkarni V, Perondi MB, Grisi S, Berg RA. A prospective investigation into the epidemiology of in-hospital pediatric cardiopulmonary resuscitation using the international Utstein reporting style. *Pediatrics* 2002;109:200–9.
471. Young KD, Gausche-Hill M, McClung CD, Lewis RJ. A prospective, population-based study of the epidemiology and outcome of out-of-hospital pediatric cardiopulmonary arrest. *Pediatrics* 2004;114:157–64.
472. Richman PB, Nashed AH. The etiology of cardiac arrest in children and young adults: special considerations for ED management. *Am J Emerg Med* 1999;17:264–70.
473. Engdahl J, Bang A, Karlson BW, Lindqvist J, Herlitz J. Characteristics and outcome among patients suffering from out-of-hospital cardiac arrest of noncardiac aetiology. *Resuscitation* 2003;57:33–41.
474. Tibballs J, Kinney S. Reduction of hospital mortality and of preventable cardiac arrest and death on introduction of a pediatric medical emergency team. *Pediatr Crit Care Med* 2009;10:306–12.
475. Hunt EA, Zimmer KP, Rinke ML, et al. Transition from a traditional code team to a medical emergency team and categorization of cardiopulmonary arrests in a children's center. *Arch Pediatr Adolesc Med* 2008;162:117–22.
476. Sharek PJ, Parast LM, Leong K, et al. Effect of a rapid response team on hospitalwide mortality and code rates outside the ICU in a Children's Hospital. *JAMA* 2007;298:2267–74.
477. Brill RJ, Gibson R, Luria JW, et al. Implementation of a medical emergency team in a large pediatric teaching hospital prevents respiratory and cardiopulmonary arrests outside the intensive care unit. *Pediatr Crit Care Med* 2007;8:236–46 [quiz 47].
478. Tibballs J, Kinney S, Duke T, Oakley E, Hennessy M. Reduction of paediatric inpatient cardiac arrest and death with a medical emergency team: preliminary results. *Arch Dis Child* 2005;90:1148–52.
479. Sagarin MJ, Chiang V, Sakles JC, et al. Rapid sequence intubation for pediatric emergency airway management. *Pediatr Emerg Care* 2002;18:417–23.
480. Moynihan RJ, Brock-Utne JG, Archer JH, Feld LH, Kreitzman TR. The effect of cricoid pressure on preventing gastric insufflation in infants and children. *Anesthesiology* 1993;78:652–6.
481. Salem MR, Joseph NJ, Heyman HJ, Belani B, Paulissian R, Ferrara TP. Cricoid compression is effective in obliterating the esophageal lumen in the presence of a nasogastric tube. *Anesthesiology* 1985;63:443–6.
482. Walker RW, Ravi R, Haylett K. Effect of cricoid force on airway calibre in children: a bronchoscopic assessment. *Br J Anaesth* 2010;104:71–4.
483. Khine HH, Corddry DH, Kettrick RG, et al. Comparison of cuffed and uncuffed endotracheal tubes in young children during general anesthesia. *Anesthesiology* 1997;86:627–31 [discussion 27A].
484. Weiss M, Dullenkopf A, Fischer JE, Keller C, Gerber AC. Prospective randomized controlled multicentre trial of cuffed or uncuffed endotracheal tubes in small children. *Br J Anaesth* 2009;103:867–73.
485. Duracher C, Schmutz E, Martinon C, Faivre J, Carli P, Orliaguet G. Evaluation of cuffed tracheal tube size predicted using the Khine formula in children. *Paediatr Anaesth* 2008;18:113–8.
486. Dullenkopf A, Kretschmar O, Knirsch W, et al. Comparison of tracheal tube cuff diameters with internal transverse diameters of the trachea in children. *Acta Anaesthesiol Scand* 2006;50:201–5.
487. Dullenkopf A, Gerber AC, Weiss M. Fit and seal characteristics of a new paediatric tracheal tube with high volume low pressure polyurethane cuff. *Acta Anaesthesiol Scand* 2005;49:232–7.
488. Salgo B, Schmitz A, Henze G, et al. Evaluation of a new recommendation for improved cuffed tracheal tube size selection in infants and small children. *Acta Anaesthesiol Scand* 2006;50:557–61.
489. Luten RC, Wears RL, Broselow J, et al. Length-based endotracheal tube and emergency equipment in pediatrics. *Ann Emerg Med* 1992;21:900–4.
490. Deakers TW, Reynolds G, Stretton M, Newth CJ. Cuffed endotracheal tubes in pediatric intensive care. *J Pediatr* 1994;125:57–62.
491. Newth CJ, Rachman B, Patel N, Hammer J. The use of cuffed versus uncuffed endotracheal tubes in pediatric intensive care. *J Pediatr* 2004;144:333–7.
492. Dorsey DP, Bowman SM, Klein MB, Archer D, Sharar SR. Perioperative use of cuffed endotracheal tubes is advantageous in young pediatric burn patients. *Burns* 2010.
493. Mhanna MJ, Zamel YB, Tichy CM, Super DM. The “air leak” test around the endotracheal tube, as a predictor of postextubation stridor, is age dependent in children. *Crit Care Med* 2002;30:2639–43.
494. Gausche M, Lewis RJ, Stratton SJ, et al. Effect of out-of-hospital pediatric endotracheal intubation on survival and neurological outcome: a controlled clinical trial. *JAMA* 2000;283:783–90.
495. Kelly JJ, Eynon CA, Kaplan JL, de Garavilla L, Dalsey WC. Use of tube condensation as an indicator of endotracheal tube placement. *Ann Emerg Med* 1998;31:575–8.

496. Andersen KH, Hald A. Assessing the position of the tracheal tube: the reliability of different methods. *Anaesthesia* 1989;44:984–5.
497. Andersen KH, SchultzLebahn T. Oesophageal intubation can be undetected by auscultation of the chest. *Acta Anaesthesiol Scand* 1994;38:580–2.
498. Van de Louw A, Cracco C, Cerf C, et al. Accuracy of pulse oximetry in the intensive care unit. *Intensive Care Med* 2001;27:1606–13.
499. Seguin P, Le Rouzo A, Tanguy M, Guillou YM, Feuillu A, Malledant Y. Evidence for the need of bedside accuracy of pulse oximetry in an intensive care unit. *Crit Care Med* 2000;28:703–6.
500. Aufderheide TP, Sigurdsson G, Pirrallo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation* 2004;109:1960–5.
501. Aufderheide TP, Lurie KG. Death by hyperventilation: a common and lifethreatening problem during cardiopulmonary resuscitation. *Crit Care Med* 2004;32:S345–51.
502. Borke WB, Munkeby BH, Morkrid L, Thaulow E, Saugstad OD. Resuscitation with 100% O₂ does not protect the myocardium in hypoxic newborn piglets. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F156–60.
503. O’Neill JF, Deakin CD. Do we hyperventilate cardiac arrest patients? *Resuscitation* 2007;73:82–5.
504. Bhende MS, Thompson AE, Orr RA. Utility of an endtidal carbon dioxide detector during stabilization and transport of critically ill children. *Pediatrics* 1992;89:1042–4.
505. Bhende MS, LaCovey DC. Endtidal carbon dioxide monitoring in the prehospital setting. *Prehosp Emerg Care* 2001;5:208–13.
506. Ornato JP, Shipley JB, Racht EM, et al. Multicenter study of a portable, handsize, colorimetric endtidal carbon dioxide detection device. *Ann Emerg Med* 1992;21:518–23.
507. Gonzalez del Rey JA, Poirier MP, Digiulio GA. Evaluation of an ambubag valve with a self-contained, colorimetric endtidal CO₂ system in the detection of airway mishaps: an animal trial. *Pediatr Emerg Care* 2000;16: 121–3.
- J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 1273 508. Bhende MS, Karasic DG, Karasic RB. Endtidal carbon dioxide changes during cardiopulmonary resuscitation after experimental asphyxial cardiac arrest. *Am J Emerg Med* 1996;14:349–50.
509. DeBehnke DJ, Hilander SJ, Dobler DW, Wickman LL, Swart GL. The hemodynamic and arterial blood gas response to asphyxiation: a canine model of pulseless electrical activity. *Resuscitation* 1995;30:169–75.
510. Ornato JP, Garnett AR, Glauser FL. Relationship between cardiac output and the endtidal carbon dioxide tension. *Ann Emerg Med* 1990;19:1104–6.
511. Mauer D, Schneider T, Elich D, Dick W. Carbon dioxide levels during prehospital active compression–decompression versus standard cardiopulmonary resuscitation. *Resuscitation* 1998;39:67–74.
512. Kolar M, Krizmaric M, Klemen P, Grmec S. Partial pressure of endtidal carbon dioxide successful predicts cardiopulmonary resuscitation in the field: a prospective observational study. *Crit Care* 2008;12:R115.
513. Sharieff GQ, Rodarte A, Wilton N, Bleyle D. The selfinflating bulb as an airway adjunct: is it reliable in children weighing less than 20 kilograms? *Acad Emerg Med* 2003;10:303–8.
514. Sharieff GQ, Rodarte A, Wilton N, Silva PD, Bleyle D. The selfinflating bulb as an esophageal detector device in children weighing more than twenty kilograms: a comparison of two techniques. *Ann Emerg Med* 2003;41:623–9.
515. Dung NM, Day NPJ, Tam DTH, et al. Fluid replacement in dengue shock syndrome: a randomized, doubleblind comparison of four intravenous fluid regimens. *Clin Infect Dis* 1999;29:787–94.
516. Ngo NT, Cao XT, Kneen R, et al. Acute management of dengue shock syndrome: a randomized doubleblind comparison of 4 intravenous fluid regimens in the first hour. *Clin Infect Dis* 2001;32:204–13.
517. Wills BA, Nguyen MD, Ha TL, et al. Comparison of three fluid solutions for resuscitation in dengue shock syndrome. *N Engl J Med* 2005;353:877–89.
518. Upadhyay M, Singhi S, Murlidharan J, Kaur N, Majumdar S. Randomized evaluation of fluid resuscitation with crystalloid (saline) and colloid (polymer from degraded gelatin in saline) in pediatric septic shock. *Indian Pediatr* 2005;42:223–31.
519. Lillis KA, Jaffe DM. Prehospital intravenous access in children. *Ann Emerg Med* 1992;21:1430–4.
520. Kanter RK, Zimmerman JJ, Strauss RH, Stoeckel KA. Pediatric emergency intravenous access. Evaluation of a protocol. *Am J Dis Child* 1986;140:132–4.
521. Kleinman ME, Oh W, Stonestreet BS. Comparison of intravenous and endotracheal epinephrine during cardiopulmonary resuscitation in newborn piglets. *Crit Care Med* 1999;27:2748–54.
522. Roberts JR, Greenburg MI, Knaub M, Baskin SI. Comparison of the pharmacological effects of epinephrine administered by the intravenous and endotracheal routes. *JACEP* 1978;7:260–4.

523. Zaritsky A. Pediatric resuscitation pharmacology. Members of the medications in pediatric resuscitation panel. *Ann Emerg Med* 1993;22:445–55.
524. Manisterski Y, Vaknin Z, BenAbraham R, et al. Endotracheal epinephrine: a call for larger doses. *Anesth Analg* 2002;95:1037–41 [table of contents].
525. Efrati O, BenAbraham R, Barak A, et al. Endobronchial adrenaline: should it be reconsidered? Dose response and haemodynamic effect in dogs. *Resuscitation* 2003;59:117–22.
526. Saul JP, Scott WA, Brown S, et al. Intravenous amiodarone for incessant tachyarrhythmias in children: a randomized, doubleblind, antiarrhythmic drug trial. *Circulation* 2005;112:3470–7.
527. Tsung JW, Blaivas M. Feasibility of correlating the pulse check with focused pointofcare echocardiography during pediatric cardiac arrest: a case series. *Resuscitation* 2008;77:264–9.
528. Cummins RO, Graves JR, Larsen MP, et al. Outofhospital transcutaneous pacing by emergency medical technicians in patients with asystolic cardiac arrest. *N Engl J Med* 1993;328:1377–82.
529. Sreeram N, Wren C. Supraventricular tachycardia in infants: response to initial treatment. *Arch Dis Child* 1990;65:127–9.
530. Benson Jr D, Smith W, Dunnigan A, Sterba R, Gallagher J. Mechanisms of regular wide QRS tachycardia in infants and children. *Am J Cardiol* 1982;49:1778–88.
531. Ackerman MJ, Siu BL, Sturner WQ, et al. Postmortem molecular analysis of SCN5A defects in sudden infant death syndrome. *JAMA* 2001;286:2264–9.
532. Arnestad M, Crotti L, Rognum TO, et al. Prevalence of longQT syndrome gene variants in sudden infant death syndrome. *Circulation* 2007;115:361–7.
533. Cronk LB, Ye B, Kaku T, et al. Novel mechanism for sudden infant death syndrome: persistent late sodium current secondary to mutations in caveolin3. *Heart Rhythm* 2007;4:161–6.
534. Millat G, Kugener B, Chevalier P, et al. Contribution of longQT syndrome genetic variants in sudden infant death syndrome. *Pediatr Cardiol* 2009;30:502–9.
535. Otagiri T, Kijima K, Osawa M, et al. Cardiac ion channel gene mutations in sudden infant death syndrome. *Pediatr Res* 2008;64:482–7.
536. Plant LD, Bowers PN, Liu Q, et al. A common cardiac sodium channel variant associated with sudden infant death in African Americans. SCN5A S1103Y. *J Clin Invest* 2006;116:430–5.
537. Tester DJ, Dura M, Carturan E, et al. A mechanism for sudden infant death syndrome (SIDS): stressinduced leak via ryanodine receptors. *Heart Rhythm* 2007;4:733–9.
538. Albert CM, Nam EG, Rimm EB, et al. Cardiac sodium channel gene variants and sudden cardiac death in women. *Circulation* 2008;117:16–23.
539. Chugh SS, Senashova O, Watts A, et al. Postmortem molecular screening in unexplained sudden death. *J Am Coll Cardiol* 2004;43:1625–9.
540. Tester DJ, Spoon DB, Valdivia HH, Makielski JC, Ackerman MJ. Targeted mutational analysis of the RyR2encoded cardiac ryanodine receptor in sudden unexplained death: a molecular autopsy of 49 medical examiner/coroner's cases. *Mayo Clin Proc* 2004;79:1380–4.
541. Graham EM, Forbus GA, Bradley SM, Shirali GS, Atz AM. Incidence and outcome of cardiopulmonary resuscitation in patients with shunted single ventricle: advantage of right ventricle to pulmonary artery shunt. *J Thorac Cardiovasc Surg* 2006;131:e7–8.
542. Charpie JR, Dekeon MK, Goldberg CS, Mosca RS, Bove EL, Kulik TJ. Postoperative hemodynamics after Norwood palliation for hypoplastic left heart syndrome. *Am J Cardiol* 2001;87:198–202.
543. Hoffman GM, Mussatto KA, Brosig CL, et al. Systemic venous oxygen saturation after the Norwood procedure and childhood neurodevelopmental outcome. *J Thorac Cardiovasc Surg* 2005;130:1094–100.
544. Johnson BA, Hoffman GM, Tweddell JS, et al. Nearinfrared spectroscopy in neonates before palliation of hypoplastic left heart syndrome. *Ann Thorac Surg* 2009;87:571–7 [discussion 7–9].
545. Hoffman GM, Tweddell JS, Ghanayem NS, et al. Alteration of the critical arteriovenous oxygen saturation relationship by sustained afterload reduction after the Norwood procedure. *J Thorac Cardiovasc Surg* 2004;127:738–45.
546. De Oliveira NC, Van Arsdell GS. Practical use of alpha blockade strategy in the management of hypoplastic left heart syndrome following stage one palliation with a BlalockTaussig shunt. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu* 2004;7:11–5.
547. Tweddell JS, Hoffman GM, Mussatto KA, et al. Improved survival of patients undergoing palliation of hypoplastic left heart syndrome: lessons learned from 115 consecutive patients. *Circulation* 2002;106:182–9.
548. Shekerdemian LS, Shore DF, Lincoln C, Bush A, Redington AN. Negativepressure ventilation improves cardiac output after right heart surgery. *Circulation* 1996;94:1149–55.

549. Shekerdemian LS, Bush A, Shore DF, Lincoln C, Redington AN. Cardiopulmonary interactions after Fontan operations: augmentation of cardiac output using negative pressure ventilation. *Circulation* 1997;96:3934–42.
550. Booth KL, Roth SJ, Thiagarajan RR, Almodovar MC, del Nido PJ, Laussen PC. Extracorporeal membrane oxygenation support of the Fontan and bidirectional Glenn circulations. *Ann Thorac Surg* 2004;77:1341–8.
551. Polderman FN, Cohen J, Blom NA, et al. Sudden unexpected death in children with a previously diagnosed cardiovascular disorder. *Int J Cardiol* 2004;95:171–6.
552. Sanatani S, Wilson G, Smith CR, Hamilton RM, Williams WG, Adatia I. Sudden unexpected death in children with heart disease. *Congenit Heart Dis* 2006;1:89–97.
553. Morris K, Beghetti M, Petros A, Adatia I, Bohn D. Comparison of hyperventilation and inhaled nitric oxide for pulmonary hypertension after repair of congenital heart disease. *Crit Care Med* 2000;28:2974–8.
554. Hoepfer MM, Galie N, Murali S, et al. Outcome after cardiopulmonary resuscitation in patients with pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2002;165:341–4.
555. Rimensberger PC, SpahrSchopfer I, Berner M, et al. Inhaled nitric oxide versus aerosolized iloprost in secondary pulmonary hypertension in children with congenital heart disease: vasodilator capacity and cellular mechanisms. *Circulation* 2001;103:544–8.
556. Liu KS, Tsai FC, Huang YK, et al. Extracorporeal life support: a simple and effective weapon for postcardiotomy right ventricular failure. *Artif Organs* 2009;33:504–8.
557. Dhillon R, Pearson GA, Firmin RK, Chan KC, Leanage R. Extracorporeal membrane oxygenation and the treatment of critical pulmonary hypertension in congenital heart disease. *Eur J Cardiothorac Surg* 1995;9:553–6.
558. Arpesella G, Loforte A, Mikus E, Mikus PM. Extracorporeal membrane oxygenation for primary allograft failure. *Transplant Proc* 2008;40:3596–7.
559. Strueber M, Hoepfer MM, Fischer S, et al. Bridge to thoracic organ transplantation in patients with pulmonary arterial hypertension using a pumpless lung assist device. *Am J Transplant* 2009;9:853–7.
560. Gluckman PD, Wyatt JS, Azzopardi D, et al. Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicentre randomised trial. *Lancet* 2005;365:663–70.
561. Battin MR, Penrice J, Gunn TR, Gunn AJ. Treatment of term infants with head cooling and mild systemic hypothermia (35.0 degrees C and 34.5 degrees C) after perinatal asphyxia. *Pediatrics* 2003;111:244–51.
562. Compagnoni G, Pogliani L, Lista G, Castoldi F, Fontana P, Mosca F. Hypothermia reduces neurological damage in asphyxiated newborn infants. *Biol Neonate* 2002;82:222–7.
563. Gunn AJ, Gunn TR, Gunning MI, Williams CE, Gluckman PD. Neuroprotection with prolonged head cooling started before postischemic seizures in fetal sheep. *Pediatrics* 1998;102:1098–106.
564. Debillon T, Daoud P, Durand P, et al. Wholebody cooling after perinatal asphyxia: a pilot study in term neonates. *Dev Med Child Neurol* 2003;45: 17–23.
565. Shankaran S, Laptook AR, Ehrenkranz RA, et al. Wholebody hypothermia for neonates with hypoxicischemic encephalopathy. *N Engl J Med* 2005;353:1574–84.
566. Doherty DR, Parshuram CS, Gaboury I, et al. Hypothermia therapy after pediatric cardiac arrest. *Circulation* 2009;119:1492–500.
567. Coimbra C, BorisMoller F, Drake M, Wieloch T. Diminished neuronal damage in the rat brain by late treatment with the antipyretic drug dipyrone or cooling following cerebral ischemia. *Acta Neuropathol* 1996;92:447–53.
568. Coimbra C, Drake M, BorisMoller F, Wieloch T. Longlasting neuroprotective effect of postischemic hypothermia and treatment with an 1274 *J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* antiinflammatory/ antipyretic drug. Evidence for chronic encephalopathic processes following ischemia. *Stroke* 1996;27:1578–85.
569. Losert H, Sterz F, Roine RO, et al. Strict normoglycaemic blood glucose levels in the therapeutic management of patients within 12 h after cardiac arrest might not be necessary. *Resuscitation* 2008;76:214–20.
570. Oksanen T, Skrifvars MB, Varpula T, et al. Strict versus moderate glucose control after resuscitation from ventricular fibrillation. *Intensive Care Med* 2007;33:2093–100.
571. PalmeKilander C. Methods of resuscitation in lowApgarscore newborn infants—a national survey. *Acta Paediatr* 1992;81:739–44.
572. Dahm LS, James LS. Newborn temperature and calculated heat loss in the delivery room. *Pediatrics* 1972;49:504–13.
573. Stephenson J, Du J, Oliver TK. The effect of cooling on blood gas tensions in newborn infants. *J Pediatr* 1970;76:848–52.

574. Gandy GM, Adamsons Jr K, Cunningham N, Silverman WA, James LS. Thermal environment and acid–base homeostasis in human infants during the first few hours of life. *J Clin Invest* 1964;43:751–8.
575. Kent AL, Williams J. Increasing ambient operating theatre temperature and wrapping in polyethylene improves admission temperature in premature infants. *J Paediatr Child Health* 2008;44:325–31.
576. Knobel RB, Wimmer Jr JE, Holbert D. Heat loss prevention for preterm infants in the delivery room. *J Perinatol* 2005;25:304–8.
577. Apgar V. A proposal for a new method of evaluation of the newborn infant. *Curr Res Anesth Analg* 1953:32.
578. Chamberlain G, Banks J. Assessment of the Apgar score. *Lancet* 1974;2:1225–8.
579. Owen CJ, Wyllie JP. Determination of heart rate in the baby at birth. *Resuscitation* 2004;60:213–7.
580. Kamlin CO, Dawson JA, O'Donnell CP, et al. Accuracy of pulse oximetry measurement of heart rate of newborn infants in the delivery room. *J Pediatr* 2008;152:756–60.
581. O'Donnell CP, Kamlin CO, Davis PG, Carlin JB, Morley CJ. Clinical assessment of infant colour at delivery. *Arch Dis Child Fetal Neonatal Ed* 2007;92:F465–7.
582. Cordero Jr L, Hon EH. Neonatal bradycardia following nasopharyngeal stimulation. *J Pediatr* 1971;78:441–7.
583. Hourii PK, Frank LR, Menegazzi JJ, Taylor R. A randomized, controlled trial of twothumb vs twofinger chest compression in a swine infant model of cardiac arrest [see comment]. *Prehosp Emerg Care* 1997;1:65–7.
584. David R. Closed chest cardiac massage in the newborn infant. *Pediatrics* 1988;81:552–4.
585. Menegazzi JJ, Auble TE, Nicklas KA, Hosack GM, Rack L, Goode JS. Twothumb versus twofinger chest compression during CRP in a swine infant model of cardiac arrest. *Ann Emerg Med* 1993;22:240–3.
586. Thaler MM, Stobie GH. An improved technique of external cardiac compression in infants and young children. *N Engl J Med* 1963;269:606–10.
587. Meyer A, Nadkarni V, Pollock A, et al. Evaluation of the Neonatal Resuscitation Program's recommended chest compression depth using computerized tomography imaging. *Resuscitation* 2010;81:544–8.
588. Wyckoff MH, Perlman JM, Laptook AR. Use of volume expansion during delivery room resuscitation in nearterm and term infants. *Pediatrics* 2005;115:950–5.
589. Soar J, Deakin CD, Nolan JP, et al. European Resuscitation Council guidelines for resuscitation 2005. Section 7. Cardiac arrest in special circumstances. *Resuscitation* 2005;67(Suppl. 1):S135–70.
590. Bronstein AC, Spyker DA, Cantilena Jr LR, Green JL, Rumack BH, Giffin SL. 2008 Annual Report of the American Association of Poison Control Centers' National Poison Data System (NPDS): 26th Annual Report. *Clin Toxicol (Phila)* 2009;47:911–1084.
591. Yanagawa Y, Sakamoto T, Okada Y. Recovery from a psychotropic drug overdose tends to depend on the time from ingestion to arrival, the Glasgow Coma Scale, and a sign of circulatory insufficiency on arrival. *Am J Emerg Med* 2007;25:757–61.
592. Zimmerman JL. Poisonings and overdoses in the intensive care unit: general and specific management issues. *Crit Care Med* 2003;31:2794–801.
593. Warner DS, Bierens JJ, Beerman SB, Katz LM. Drowning: a cry for help. *Anesthesiology* 2009;110:1211–3.
594. Danzl D. Accidental hypothermia. In: Auerbach P, editor. *Wilderness medicine*. St. Louis: Mosby; 2007. p. 125–60.
595. Paal P, Beikircher W, Brugger H. Avalanche emergencies. Review of the current situation. *Anaesthesist* 2006;55:314–24.
596. Mattu A, Brady WJ, Perron AD. Electrocardiographic manifestations of hypothermia. *Am J Emerg Med* 2002;20:314–26.
597. Ujhelyi MR, Sims JJ, Dubin SA, Vender J, Miller AW. Defibrillation energy requirements and electrical heterogeneity during total body hypothermia. *Crit Care Med* 2001;29:1006–11.
598. Walpoth BH, Walpoth Aslan BN, Mattle HP, et al. Outcome of survivors of accidental deep hypothermia and circulatory arrest treated with extracorporeal blood warming. *N Engl J Med* 1997;337:1500–5.
599. Bouchama A, Knochel JP. Heat stroke. *N Engl J Med* 2002;346:1978–88.
600. Bouchama A. The 2003 European heat wave. *Intensive Care Med* 2004;30:1–3.
601. Coris EE, Ramirez AM, Van Durme DJ. Heat illness in athletes: the dangerous combination of heat, humidity and exercise. *Sports Med* 2004;34:9–16.
602. Grogan H, Hopkins PM. Heat stroke: implications for critical care and anaesthesia. *Br J Anaesth* 2002;88:700–7.

603. Hadad E, Weinbroum AA, BenAbraham R. Druginduced hyperthermia and muscle rigidity: a practical approach. *Eur J Emerg Med* 2003;10:149–54.
604. Halloran LL, Bernard DW. Management of druginduced hyperthermia. *Curr Opin Pediatr* 2004;16:211–5.
605. Bouchama A, Dehbi M, ChavesCarballo E. Cooling and hemodynamic management in heatstroke: practical recommendations. *Crit Care* 2007;11:R54.
606. Eshel G, Safar P, Sassano J, Stezoski W. Hyperthermia-induced cardiac arrest in dogs and monkeys. *Resuscitation* 1990;20:129–43.
607. Eshel G, Safar P, Radovsky A, Stezoski SW. Hyperthermia-induced cardiac arrest in monkeys: limited efficacy of standard CPR. *Aviat Space Environ Med* 1997;68:415–20.
608. Masoli M, Fabian D, Holt S, Beasley R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004;59:469–78.
609. Global Strategy for Asthma Management and Prevention 2009 [accessed 24.06.10].
610. Williams TJ, Tuxen DV, Scheinkestel CD, Czarny D, Bowes G. Risk factors for morbidity in mechanically ventilated patients with acute severe asthma. *Am Rev Respir Dis* 1992;146:607–15.
611. Bowman FP, Menegazzi JJ, Check BD, Duckett TM. Lower esophageal sphincter pressure during prolonged cardiac arrest and resuscitation. *Ann Emerg Med* 1995;26:216–9.
612. Leatherman JW, McArthur C, Shapiro RS. Effect of prolongation of expiratory time on dynamic hyperinflation in mechanically ventilated patients with severe asthma. *Crit Care Med* 2004;32:1542–5.
613. Lapinsky SE, Leung RS. AutoPEEP and electromechanical dissociation. *N Engl J Med* 1996;335:674.
614. Rogers PL, Schlichtig R, Miro A, Pinsky M. AutoPEEP during CPR. An “occult” cause of electromechanical dissociation? *Chest* 1991;99:492–3.
615. Rosengarten PL, Tuxen DV, Dziukas L, Scheinkestel C, Merrett K, Bowes G. Circulatory arrest induced by intermittent positive pressure ventilation in a patient with severe asthma. *Anaesth Intensive Care* 1991;19:118–21.
616. Sprung J, Hunter K, Barnas GM, Bourke DL. Abdominal distention is not always a sign of esophageal intubation: cardiac arrest due to “autoPEEP”. *Anesth Analg* 1994;78:801–4.
617. Harrison R. Chest compression first aid for respiratory arrest due to acute asphyxic asthma. *Emerg Med J* 2010;27:59–61.
618. Deakin CD, McLaren RM, Petley GW, Clewlow F, DalrympleHay MJ. Effects of positive endexpiratory pressure on transthoracic impedance—implications for defibrillation. *Resuscitation* 1998;37:9–12.
619. Galbois A, AitOufella H, Baudel JL, et al. Pleural ultrasound compared to chest radiographic detection of pneumothorax resolution after drainage. *Chest* 2010.
620. Mabuchi N, Takasu H, Ito S, et al. Successful extracorporeal lung assist (ECLA) for a patient with severe asthma and cardiac arrest. *Clin Intensive Care* 1991;2:292–4.
621. Martin GB, Rivers EP, Paradis NA, Goetting MG, Morris DC, Nowak RM. Emergency department cardiopulmonary bypass in the treatment of human cardiac arrest. *Chest* 1998;113:743–51.
622. Soar J, Pumphrey R, Cant A, et al. Emergency treatment of anaphylactic reactions—guidelines for healthcare providers. *Resuscitation* 2008;77: 157–69.
623. Soar J. Emergency treatment of anaphylaxis in adults: concise guidance. *Clin Med* 2009;9:181–5.
624. Charalambous CP, Zipitis CS, Keenan DJ. Chest reexploration in the intensive care unit after cardiac surgery: a safe alternative to returning to the operating theater. *Ann Thorac Surg* 2006;81:191–4.
625. McKowen RL, Magovern GJ, Liebler GA, Park SB, Burkholder JA, Maher TD. Infectious complications and costeffectiveness of open resuscitation in the surgical intensive care unit after cardiac surgery. *Ann Thorac Surg* 1985;40:388–92.
626. Pottle A, Bullock I, Thomas J, Scott L. Survival to discharge following Open Chest Cardiac Compression (OCCC). A 4year retrospective audit in a cardiothoracic specialist centre – Royal Brompton and Harefield NHS Trust, United Kingdom. *Resuscitation* 2002;52:269–72.
627. Mackay JH, Powell SJ, Osgathorp J, Rozario CJ. Sixyear prospective audit of chest reopening after cardiac arrest. *Eur J Cardiothorac Surg* 2002;22:421–5.
628. Birdi I, Chaudhuri N, Lenthall K, Reddy S, Nashef SA. Emergency reinstatement of cardiopulmonary bypass following cardiac surgery: outcome justifies the cost. *Eur J Cardiothorac Surg* 2000;17:743–6.
629. elBanayosy A, Brehm C, Kizner L, et al. Cardiopulmonary resuscitation after cardiac surgery: a twoyear study. *J Cardiothorac Vasc Anesth* 1998;12:390–2.
630. Anthi A, Tzelepis GE, Alivizatos P, Michalis A, Palatianos GM, Geroulanos S. Unexpected cardiac arrest after cardiac surgery: incidence, predisposing causes, and outcome of open chest cardiopulmonary resuscitation. *Chest* 1998;113:15–9.

631. Wahba A, Gotz W, Birnbaum DE. Outcome of cardiopulmonary resuscitation following open heart surgery. *Scand Cardiovasc J* 1997;31:147–9.
632. Kaiser GC, Naunheim KS, Fiore AC, et al. Reoperation in the intensive care unit. *Ann Thorac Surg* 1990;49:903–7 [discussion 8].
633. Rhodes JF, Blaufox AD, Seiden HS, et al. Cardiac arrest in infants after congenital heart surgery. *Circulation* 1999;100:II194–9.
634. Kempen PM, Allgood R. Right ventricular rupture during closedchest cardiopulmonary resuscitation after pneumonectomy with pericardiotomy: a case report. *Crit Care Med* 1999;27:1378–9.
635. Bohrer H, Gust R, Bottiger BW. Cardiopulmonary resuscitation after cardiac surgery. *J Cardiothorac Vasc Anesth* 1995;9:352.
636. Klintschar M, Darok M, Radner H. Massive injury to the heart after attempted active compression–decompression cardiopulmonary resuscitation. *Int J Legal Med* 1998;111:93–6.
637. Fosse E, Lindberg H. Left ventricular rupture following external chest compression. *Acta Anaesthesiol Scand* 1996;40:502–4.
- J.P. Nolan et al. / Resuscitation 81 (2010) 1219–1276* 1275
638. Li Y, Wang H, Cho JH, et al. Defibrillation delivered during the upstroke phase of manual chest compression improves shock success. *Crit Care Med* 2010;38:910–5.
639. Li Y, Yu T, Ristagno G, et al. The optimal phasic relationship between synchronized shock and mechanical chest compressions. *Resuscitation* 2010;81:724–9.
640. Rosemurgy AS, Norris PA, Olson SM, Hurst JM, Albrink MH. Prehospital traumatic cardiac arrest: the cost of futility. *J Trauma* 1993;35:468–73.
641. Shimazu S, Shatney CH. Outcomes of trauma patients with no vital signs on hospital admission. *J Trauma* 1983;23:213–6.
642. Battistella FD, Nugent W, Owings JT, Anderson JT. Field triage of the pulseless trauma patient. *Arch Surg* 1999;134:742–5.
643. Stockinger ZT, McSwain Jr NE. Additional evidence in support of withholding or terminating cardiopulmonary resuscitation for trauma patients in the field. *J Am Coll Surg* 2004;198:227–31.
644. Fulton RL, Voigt WJ, Hilakos AS. Confusion surrounding the treatment of traumatic cardiac arrest. *J Am Coll Surg* 1995;181:209–14.
645. Pasquale MD, Rhodes M, Cipolle MD, Hanley T, Wasser T. Defining “dead on arrival”: impact on a level I trauma center. *J Trauma* 1996;41:726–30.
646. Stratton SJ, Brickett K, Crammer T. Prehospital pulseless, unconscious penetrating trauma victims: field assessments associated with survival. *J Trauma* 1998;45:96–100.
647. Maron BJ, Estes 3rd NA. Commotio cordis. *N Engl J Med* 2010;362:917–27.
648. Maron BJ, Gohman TE, Kyle SB, Estes 3rd NA, Link MS. Clinical profile and spectrum of commotio cordis. *JAMA* 2002;287:1142–6.
649. Maron BJ, Estes 3rd NA, Link MS. Task Force 11: commotio cordis. *J Am Coll Cardiol* 2005;45:1371–3.
650. Nesbitt AD, Cooper PJ, Kohl P. Rediscovering commotio cordis. *Lancet* 2001;357:1195–7.
651. Link MS, Estes M, Maron BJ. Sudden death caused by chest wall trauma (commotio cordis). In: Kohl P, Sachs F, Franz MR, editors. *Cardiac mechanoelectric feedback and arrhythmias: from pipette to patient*. Philadelphia: Elsevier Saunders; 2005. p. 270–6.
652. Cera SM, Mostafa G, Sing RF, Sarafin JL, Matthews BD, Heniford BT. Physiologic predictors of survival in posttraumatic arrest. *Am Surg* 2003;69:140–4.
653. Esposito TJ, Jurkovich GJ, Rice CL, Maier RV, Copass MK, Ashbaugh DG. Reappraisal of emergency room thoracotomy in a changing environment. *J Trauma* 1991;31:881–5 [discussion 5–7].
654. Martin SK, Shatney CH, Sherck JP, et al. Blunt trauma patients with prehospital pulseless electrical activity (PEA): poor ending assured. *J Trauma* 2002;53:876–80 [discussion 80–81].
655. Domeier RM, McSwain Jr NE, Hopson LR, et al. Guidelines for withholding or termination of resuscitation in prehospital traumatic cardiopulmonary arrest. *J Am Coll Surg* 2003;196:475–81.
656. Gervin AS, Fischer RP. The importance of prompt transport of salvage of patients with penetrating heart wounds. *J Trauma* 1982;22:443–8.
657. Branney SW, Moore EE, Feldhaus KM, Wolfe RE. Critical analysis of two decades of experience with postinjury emergency department thoracotomy in a regional trauma center. *J Trauma* 1998;45:87–94 [discussion 5].
658. Durham III LA, Richardson RJ, Wall Jr MJ, Pepe PE, Mattox KL. Emergency center thoracotomy: impact of prehospital resuscitation. *J Trauma* 1992;32:775–9.
659. Frezza EE, Mezgebe H. Is 30 minutes the golden period to perform emergency room thoracotomy (ERT) in penetrating chest injuries? *J Cardiovasc Surg (Torino)* 1999;40:147–51.

660. Powell DW, Moore EE, Cothren CC, et al. Is emergency department resuscitative thoracotomy futile care for the critically injured patient requiring prehospital cardiopulmonary resuscitation? *J Am Coll Surg* 2004;199:211–5.
661. Coats TJ, Keogh S, Clark H, Neal M. Prehospital resuscitative thoracotomy for cardiac arrest after penetrating trauma: rationale and case series. *J Trauma* 2001;50:670–3.
662. Wise D, Davies G, Coats T, Lockey D, Hyde J, Good A. Emergency thoracotomy: “how to do it”. *Emerg Med J* 2005;22:22–4.
663. Kwan I, Bunn F, Roberts I. Spinal immobilisation for trauma patients. *Cochrane Database Syst Rev* 2001:CD002803.
664. Practice management guidelines for emergency department thoracotomy. Working Group, Ad Hoc Subcommittee on Outcomes, American College of Surgeons Committee on Trauma. *J Am Coll Surg* 2001;193:303–9.
665. Walcher F, Kortum S, Kirschning T, Weihgold N, Marzi I. Optimized management of polytraumatized patients by prehospital ultrasound. *Unfallchirurg* 2002;105:986–94.
666. Kirschning T, Brenner F, Stier M, Weber CF, Walcher F. Prehospital emergency sonography of trauma patients. *Anaesthesist* 2009;58:51–60.
667. Department of Health, Welsh Office, Scottish Office Department of Health, Department of Health and Social Services, Northern Ireland. Why mothers die. Report on confidential enquiries into maternal deaths in the United Kingdom, 2000–2002. London: The Stationery Office; 2004.
668. Hogan MC, Foreman KJ, Naghavi M, et al. Maternal mortality for 181 countries, 1980–2008: a systematic analysis of progress towards Millennium Development Goal 5. *Lancet* 2010;375:1609–23.
669. Lewis G. The Confidential Enquiry into Maternal and Child Health (CEMACH). Saving Mothers’ Lives: reviewing maternal deaths to make motherhood safer – 2003–2005. The Seventh Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. London: CEMACH; 2007.
670. PageRodriguez A, GonzalezSanchez JA. Perimortem cesarean section of twin pregnancy: case report and review of the literature. *Acad Emerg Med* 1999;6:1072–4.
671. Cardosi RJ, Porter KB. Cesarean delivery of twins during maternal cardiopulmonary arrest. *Obstet Gynecol* 1998;92:695–7.
672. Johnson MD, Luppi CJ, Over DC. Cardiopulmonary resuscitation. In: Gambling DR, Douglas MJ, editors. *Obstetric anesthesia and uncommon disorders*. Philadelphia: W.B. Saunders; 1998. p. 51–74.
673. Nanson J, Elcock D, Williams M, Deakin CD. Do physiological changes in pregnancy change defibrillation energy requirements? *Br J Anaesth* 2001;87:237–9.
674. Katz VL, Dotters DJ, Droegemueller W. Perimortem cesarean delivery. *Obstet Gynecol* 1986;68:571–6.
675. American Heart Association in collaboration with International Liaison Committee on Resuscitation. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2000;102(Suppl.):11–384.
676. Chapter 4; Part 6: cardiac arrest associated with pregnancy. Cummins R, Hazinski M, Field J, editors. *ACLS—the reference textbook*. Dallas: American Heart Association; 2003. p. 143–58.
677. Budnick LD. Bathtub-related electrocutions in the United States, 1979 to 1982. *JAMA* 1984;252:918–20.
678. Lightning-associated deaths—United States, 1980–1995. *MMWR Morb Mortal Wkly Rep* 1998;47:391–4.
679. Geddes LA, Bourland JD, Ford G. The mechanism underlying sudden death from electric shock. *Med Instrum* 1986;20:303–15.
680. Cooper MA. Lightning injuries: prognostic signs for death. *Ann Emerg Med* 1980;9:134–8.
681. KleinschmidtDeMasters BK. Neuropathology of lightningstrike injuries. *Semin Neurol* 1995;15:323–8.
682. Stewart CE. When lightning strikes. *EmergMedServ* 2000;29:57–67 [quiz 103].
683. Cooper MA. Emergent care of lightning and electrical injuries. *Semin Neurol* 1995;15:268–78.
684. Duclos PJ, Sanderson LM. An epidemiological description of lightning-related deaths in the United States. *Int J Epidemiol* 1990;19:673–9.
685. Epperly TD, Stewart JR. The physical effects of lightning injury. *J Fam Pract* 1989;29:267–72.
686. Whitcomb D, Martinez JA, Daberkow D. Lightning injuries. *South Med J* 2002;95:1331–4.
687. Chamberlain DA, Hazinski MF. Education in resuscitation. *Resuscitation* 2003;59:11–43.
688. Yeung J, Perkins GD. Timing of drug administration during CPR and the role of simulation. *Resuscitation* 2010;81:265–6.

689. Berdowski J, Schmohl A, Tijssen JG, Koster RW. Time needed for a regional emergency medical system to implement resuscitation Guidelines 2005—The Netherlands experience. *Resuscitation* 2009;80:1336–41.
690. Bigham BL, Koprowicz K, Aufderheide TP, et al. Delayed Prehospital Implementation of the 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiac Care. *Prehosp Emerg Care* 2010.
691. Andersen PO, Jensen MK, Lippert A, Ostergaard D. Identifying nontechnical skills and barriers for improvement of teamwork in cardiac arrest teams. *Resuscitation* 2010;81:695–702.
692. Flin R, Patey R, Glavin R, Maran N. Anaesthetists' nontechnical skills. *Br J Anaesth* 2010.
693. Axelsson A, Thoren A, Holmberg S, Herlitz J. Attitudes of trained Swedish lay rescuers toward CPR performance in an emergency: a survey of 1012 recently trained CPR rescuers. *Resuscitation* 2000;44:27–36.
694. Hubble MW, Bachman M, Price R, Martin N, Huie D. Willingness of high school students to perform cardiopulmonary resuscitation and automated external defibrillation. *Prehosp Emerg Care* 2003;7:219–24.
695. Swor RA, Jackson RE, Compton S, et al. Cardiac arrest in private locations: different strategies are needed to improve outcome. *Resuscitation* 2003;58:171–6.
696. Swor R, Khan I, Domeier R, Honeycutt L, Chu K, Compton S. CPR training and CPR performance: do CPR-trained bystanders perform CPR? *Acad Emerg Med* 2006;13:596–601.
697. Vaillancourt C, Stiell IG, Wells GA. Understanding and improving low bystander CPR rates: a systematic review of the literature. *CJEM* 2008;10:51–65.
698. Boucek CD, Phrampus P, Lutz J, Dongilli T, Bircher NG. Willingness to perform mouth-to-mouth ventilation by health care providers: a survey. *Resuscitation* 2009;80:849–53.
699. Caves ND, Irwin MG. Attitudes to basic life support among medical students following the 2003 SARS outbreak in Hong Kong. *Resuscitation* 2006;68:93–100.
700. Coons SJ, Guy MC. Performing bystander CPR for sudden cardiac arrest: behavioral intentions among the general adult population in Arizona. *Resuscitation* 2009;80:334–40.
701. Dwyer T. Psychological factors inhibit family members' confidence to initiate CPR. *Prehosp Emerg Care* 2008;12:157–61.
702. Jelinek GA, Gennat H, Celenza T, O'Brien D, Jacobs I, Lynch D. Community attitudes towards performing cardiopulmonary resuscitation in Western Australia. *Resuscitation* 2001;51:239–46.
703. Johnston TC, Clark MJ, Dingle GA, FitzGerald G. Factors influencing Queenslanders' willingness to perform bystander cardiopulmonary resuscitation. *Resuscitation* 2003;56:67–75.
704. Kuramoto N, Morimoto T, Kubota Y, et al. Public perception of and willingness to perform bystander CPR in Japan. *Resuscitation* 2008;79:475–81.
705. Omi W, Taniguchi T, Kaburaki T, et al. The attitudes of Japanese high school students toward cardiopulmonary resuscitation. *Resuscitation* 2008;78:340–5.
706. Riegel B, Mosesso VN, Birnbaum A, et al. Stress reactions and perceived difficulties of lay responders to a medical emergency. *Resuscitation* 2006;70:98–106.
707. Shibata K, Taniguchi T, Yoshida M, Yamamoto K. Obstacles to bystander cardiopulmonary resuscitation in Japan. *Resuscitation* 2000;44:187–93.
- 1276 J.P. Nolan et al. / *Resuscitation* 81 (2010) 1219–1276 708. Taniguchi T, Omi W, Inaba H. Attitudes toward the performance of bystander cardiopulmonary resuscitation in Japan. *Resuscitation* 2007;75:82–7.
709. Moser DK, Dracup K, Doering LV. Effect of cardiopulmonary resuscitation training for parents of high-risk neonates on perceived anxiety, control, and burden. *Heart Lung* 1999;28:326–33.
710. Axelsson A, Herlitz J, Ekstrom L, Holmberg S. Bystander-initiated cardiopulmonary resuscitation out-of-hospital. A first description of the bystanders and their experiences. *Resuscitation* 1996;33:3–11.
711. Donohoe RT, Haefeli K, Moore F. Public perceptions and experiences of myocardial infarction, cardiac arrest and CPR in London. *Resuscitation* 2006;71:70–9.
712. Hamasu S, Morimoto T, Kuramoto N, et al. Effects of BLS training on factors associated with attitude toward CPR in college students. *Resuscitation* 2009;80:359–64.
713. Parnell MM, Pearson J, Galletly DC, Larsen PD. Knowledge of and attitudes towards resuscitation in New Zealand high school students. *Emerg Med J* 2006;23:899–902.
714. Swor R, Compton S, Farr L, et al. Perceived self-efficacy in performing and willingness to learn cardiopulmonary resuscitation in an elderly population in a suburban community. *Am J Crit Care* 2003;12:65–70.
715. Perkins GD, Walker G, Christensen K, Hulme J, Monsieurs KG. Teaching recognition of agonal breathing improves accuracy of diagnosing cardiac arrest. *Resuscitation* 2006;70:432–7.
716. Yeung J, Meeks R, Edelson D, Gao F, Soar J, Perkins GD. The use of CPR feedback/prompt devices during training and CPR performance: a systematic review. *Resuscitation* 2009;80:743–51.

717. Lam KK, Lau FL, Chan WK, Wong WN. Effect of severe acute respiratory syndrome on bystander willingness to perform cardiopulmonary resuscitation (CPR)—is compression only preferred to standard CPR? *Prehosp Disaster Med* 2007;22:325–9.
718. Locke CJ, Berg RA, Sanders AB, et al. Bystander cardiopulmonary resuscitation. Concerns about mouth-to-mouth contact. *Arch Intern Med* 1995;155:938–43.
719. Hoke RS, Chamberlain DA, Handley AJ. A reference automated external defibrillator provider course for Europe. *Resuscitation* 2006;69:421–33.
720. Lynch B, Einspruch EL, Nichol G, Becker LB, Aufderheide TP, Idris A. Effectiveness of a 30min CPR self-instruction program for lay responders: a controlled randomized study. *Resuscitation* 2005;67:31–43.
721. Todd KH, Braslow A, Brennan RT, et al. Randomized, controlled trial of video self-instruction versus traditional CPR training. *Ann Emerg Med* 1998;31:364–9.
722. Einspruch EL, Lynch B, Aufderheide TP, Nichol G, Becker L. Retention of CPR skills learned in a traditional AHA Heartsaver course versus 30min video self-training: a controlled randomized study. *Resuscitation* 2007;74:476–86.
723. Todd KH, Heron SL, Thompson M, Dennis R, O'Connor J, Kellermann AL. Simple CPR: a randomized, controlled trial of video self-instructional cardiopulmonary resuscitation training in an African American church congregation. *Ann Emerg Med* 1999;34:730–7.
724. Reder S, Cummings P, Quan L. Comparison of three instructional methods for teaching cardiopulmonary resuscitation and use of an automatic external defibrillator to high school students. *Resuscitation* 2006;69:443–53.
725. Roppolo LP, Pepe PE, Campbell L, et al. Prospective, randomized trial of the effectiveness and retention of 30min layperson training for cardiopulmonary resuscitation and automated external defibrillators: The American Airlines Study. *Resuscitation* 2007;74:276–85.
726. Batcheller AM, Brennan RT, Braslow A, Urrutia A, Kaye W. Cardiopulmonary resuscitation performance of subjects over forty is better following half-hour video self-instruction compared to traditional four-hour classroom training. *Resuscitation* 2000;43:101–10.
727. Braslow A, Brennan RT, Newman MM, Bircher NG, Batcheller AM, Kaye W. CPR training without an instructor: development and evaluation of a video self-instructional system for effective performance of cardiopulmonary resuscitation. *Resuscitation* 1997;34:207–20.
728. Isbye DL, Rasmussen LS, Lippert FK, Rudolph SF, Ringsted CV. Laypersons may learn basic life support in 24 min using a personal resuscitation manikin. *Resuscitation* 2006;69:435–42.
729. Moule P, Albarran JW, Bessant E, Brownfield C, Pollock J. A nonrandomized comparison of e-learning and classroom delivery of basic life support with automated external defibrillator use: a pilot study. *Int J Nurs Pract* 2008;14:427–34.
730. Liberman M, Golberg N, Mulder D, Sampalis J. Teaching cardiopulmonary resuscitation to CEGEP students in Quebec—a pilot project. *Resuscitation* 2000;47:249–57.
731. Jones I, Handley AJ, Whitfield R, Newcombe R, Chamberlain D. A preliminary feasibility study of a short DVD-based distance learning package for basic life support. *Resuscitation* 2007;75:350–6.
732. Brannon TS, White LA, Kilcrease JN, Richard LD, Spillers JG, Phelps CL. Use of instructional video to prepare parents for learning infant cardiopulmonary resuscitation. *Proc (Bayl Univ Med Cent)* 2009;22:133–7.
733. de Vries W, Turner N, Monsieurs K, Bierens J, Koster R. Comparison of instructor-led automated external defibrillation training and three alternative DVD-based training methods. *Resuscitation* 2010;81:1004–9.
734. Perkins GD, Mancini ME. Resuscitation training for healthcare workers. *Resuscitation* 2009;80:841–2.
735. Spooner BB, Fallaha JF, Kocierz L, Smith CM, Smith SC, Perkins GD. An evaluation of objective feedback in basic life support (BLS) training. *Resuscitation* 2007;73:417–24.
736. Andresen D, Arntz HR, Grafing W, et al. Public access resuscitation program including defibrillator training for laypersons: a randomized trial to evaluate the impact of training course duration. *Resuscitation* 2008;76:419–24.
737. Smith KK, Gilcreast D, Pierce K. Evaluation of staff's retention of ACLS and BLS skills. *Resuscitation* 2008;78:59–65.
738. Woollard M, Whitfield R, Smith A, et al. Skill acquisition and retention in automated external defibrillator (AED) use and CPR by lay responders: a prospective study. *Resuscitation* 2004;60:17–28.
739. Berden HJ, Willems FF, Hendrick JM, Pijls NH, Knape JT. How frequently should basic cardiopulmonary resuscitation training be repeated to maintain adequate skills? *BMJ* 1993;306:1576–7.

740. Woollard M, Whitfield R, Newcombe RG, Colquhoun M, Vetter N, Chamberlain D. Optimal refresher training intervals for AED and CPR skills: a randomised controlled trial. *Resuscitation* 2006;71:237–47.
741. Riegel B, Nafziger SD, McBurnie MA, et al. How well are cardiopulmonary resuscitation and automated external defibrillator skills retained over time? Results from the Public Access Defibrillation (PAD) trial. *Acad Emerg Med* 2006;13:254–63.
742. Beckers SK, Fries M, Bickenbach J, et al. Retention of skills in medical students following minimal theoretical instructions on semi and fully automated external defibrillators. *Resuscitation* 2007;72:444–50.
743. Perkins GD, Lockett AS. Defibrillationsafety versus efficacy. *Resuscitation* 2008;79:1–3.
744. Perkins GD, Barrett H, Bullock I, et al. The Acute Care Undergraduate TEaching (ACUTE) Initiative: consensus development of core competencies in acute care for undergraduates in the United Kingdom. *Intensive Care Med* 2005;31:1627–33.
745. Schwid HA, Rooke GA, Ross BK, Sivarajan M. Use of a computerized advanced cardiac life support simulator improves retention of advanced cardiac life support guidelines better than a textbook review. *Crit Care Med* 1999;27:821–4.
746. Polglase RF, Parish DC, Buckley RL, Smith RW, Joiner TA. Problembased ACLS instruction: a model approach for undergraduate emergency medical education. *Ann Emerg Med* 1989;18:997–1000.
747. Clark LJ, Watson J, Cobbe SM, Reeve W, Swann IJ, Macfarlane PW. CPR '98: a practical multimedia computerbased guide to cardiopulmonary resuscitation for medical students. *Resuscitation* 2000;44:109–17.
748. Hudson JN. Computeraided learning in the real world of medical education: does the quality of interaction with the computer affect student learning? *Med Educ* 2004;38:887–95.
749. Jang KS, Hwang SY, Park SJ, Kim YM, Kim MJ. Effects of a webbased teaching method on undergraduate nursing students' learning of electrocardiography. *J Nurs Educ* 2005;44:35–9.
750. Kim JH, Kim WO, Min KT, Yang JY, Nam YT. Learning by computer simulation does not lead to better test performance than textbook study in the diagnosis and treatment of dysrhythmias. *J Clin Anesth* 2002;14:395–400.
751. Leong SL, Baldwin CD, Adelman AM. Integrating webbased computer cases into a required clerkship: development and evaluation. *Acad Med* 2003;78:295–301.
752. Rosser JC, Herman B, Risucci DA, Murayama M, Rosser LE, Merrell RC. Effectiveness of a CDROM multimedia tutorial in transferring cognitive knowledge essential for laparoscopic skill training. *Am J Surg* 2000;179:320–4.
753. Papadimitriou L, Xanthos T, Bassiakou E, Stroumpoulis K, Barouxis D, Iacovidou N. Distribution of precourse BLS/AED manuals does not influence skill acquisition and retention in lay rescuers: a randomised study. *Resuscitation* 2010;81:348–52.
754. Perkins GD. Simulation in resuscitation training. *Resuscitation* 2007;73:202–11.
755. Duran R, Aladag N, Vatansever U, Kucukugurluoglu Y, Sut N, Acunas B. Proficiency and knowledge gained and retained by pediatric residents after neonatal resuscitation course. *Pediatr Int* 2008;50:644–7.
756. Anthonypillai F. Retention of advanced cardiopulmonary resuscitation knowledge by intensive care trained nurses. *Intensive Crit Care Nurs* 1992;8:180–4.
757. Boonmak P, Boonmak S, Srichaipanha S, Poomsawat S. Knowledge and skill after brief ACLS training. *J Med Assoc Thai* 2004;87:1311–4.
758. Kaye W, Wynne G, Marteau T, et al. An advanced resuscitation training course for preregistration house officers. *J R Coll Physicians Lond* 1990;24:51–4.
759. Semeraro F, Signore L, Cerchiari EL. Retention of CPR performance in anaesthetists. *Resuscitation* 2006;68:101–8.
760. Skidmore MB, Urquhart H. Retention of skills in neonatal resuscitation. *Paediatr Child Health* 2001;6:31–5.
761. Trevisanuto D, Ferrarese P, Cavicchioli P, Fasson A, Zanardo V, Zacchello F. Knowledge gained by pediatric residents after neonatal resuscitation program courses. *Paediatr Anaesth* 2005;15:944–7.
762. Young R, King L. An evaluation of knowledge and skill retention following an inhouse advanced life support course. *Nurs Crit Care* 2000;5:7–14.
763. Grant EC, Marczynski CA, Menon K. Using pediatric advanced life support in pediatric residency training: does the curriculum need resuscitation? *Pediatr Crit Care Med* 2007;8:433–9.
764. O'Steen DS, Kee CC, Minick MP. The retention of advanced cardiac life support knowledge among registered nurses. *J Nurs Staff Dev* 1996;12:66–72.
765. Hammond F, Saba M, Simes T, Cross R. Advanced life support: retention of registered nurses' knowledge 18 months after initial training. *Aust Crit Care* 2000;13:99–104.
766. Baskett PJ, Lim A. The varying ethical attitudes towards resuscitation in Europe. *Resuscitation* 2004;62:267–73.