

Odporúčania Európskej resuscitačnej rady pre kardiopulmonálnu resuscitáciu 2010

Sekcia 6 Resuscitácia u detí

Resuscitation 81 (2010) 1364–1388

Zoznam literatúry

1. European Resuscitation Council. Paediatric life support: (including the recommendations for resuscitation of babies at birth). *Resuscitation* 1998;37:95–6.
2. Zideman D, Bingham R, Beattie T, et al. Guidelines for paediatric life support: a statement by the Paediatric Life Support Working Party of the European Resuscitation Council, 1993. *Resuscitation* 1994;27:91–105.
3. Phillips B, Zideman D, Wyllie J, Richmond S, van Reempts P. European Resuscitation Council Guidelines 2000 for newly born life support. A statement from the Paediatric Life Support Working Group and approved by the Executive Committee of the European Resuscitation Council. *Resuscitation* 2001;48:235–9.
4. Phillips B, Zideman D, GarciaCastrillo L, Felix M, ShwarzSchwierin V. European Resuscitation Council Guidelines 2000 for advanced paediatric life support. A statement from Paediatric Life Support Working Group and approved by the Executive Committee of the European Resuscitation Council. *Resuscitation* 2001;48:231–4.
5. Biarent D, Bingham R, Richmond S, et al. European Resuscitation Council Guidelines for Resuscitation 2005. Section 6. Paediatric life support. *Resuscitation* 2005;67:S97–133.
6. American Heart Association in collaboration with International Liaison Committee on Resuscitation. Guidelines for cardiopulmonary resuscitation and emergency cardiovascular care – an international consensus on science. *Resuscitation* 2000;46:3–430.
7. American Heart Association in collaboration with International Liaison Committee on Resuscitation. Guidelines 2000 for cardiopulmonary resuscitation and emergency cardiovascular care: international consensus on science. *Circulation* 2000;102(Suppl. I):I46–8.
8. 2005 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. Part 6: Paediatric basic and advanced life support. *Resuscitation* 2005;67: 271–91.
9. 2010 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation*, 2010; in press.
10. 2010 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation*, 2010; in press.
11. Richmond S, Wyllie J. European Resuscitation Council Guidelines for Resuscitation 2010. Section 7. Resuscitation of babies at birth. *Resuscitation* 2010;81:1389–99.
12. Tibballs J, Weeraratna C. The influence of time on the accuracy of healthcare personnel to diagnose paediatric cardiac arrest by pulse palpation. *Resuscitation* 2010;81:671–5.
13. Tibballs J, Carter B, Kiraly NJ, Ragg P, Clifford M. External and internal biphasic direct current shock doses for pediatric ventricular fibrillation and pulseless ventricular tachycardia. *Pediatr Crit Care Med* 2010.
14. Sarti A, Savron F, Ronfani L, Pelizzo G, Barbi E. Comparison of three sites to check the pulse and count heart rate in hypotensive infants. *Paediatr Anaesth* 2006;16:394–8.
15. Sarti A, Savron F, Casotto V, Cuttini M. Heartbeat assessment in infants: a comparison of four clinical methods. *Pediatr Crit Care Med* 2005;6:212–5.
16. de Caen AR, Kleinman ME, Chameides L, et al. International consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. Part 10: Pediatric basic and advanced life support.
17. Berg RA, Hilwig RW, Kern KB, Babar I, Ewy GA. Simulated mouth-to-mouth ventilation and chest compressions (bystander cardiopulmonary resuscitation) improves outcome in a swine model of prehospital pediatric asphyxial cardiac arrest. *Crit Care Med* 1999;27:1893–9.
18. Dorph E, Wik L, Steen PA. Effectiveness of ventilation-compression ratios 1:5 and 2:15 in simulated single rescuer paediatric resuscitation. *Resuscitation* 2002;54:259–64.
19. Turner I, Turner S, Armstrong V. Does the compression to ventilation ratio affect the quality of CPR: a simulation study. *Resuscitation* 2002;52:55–62.

20. Babbs CF, Kern KB. Optimum compression to ventilation ratios in CPR under realistic, practical conditions: a physiological and mathematical analysis. *Resuscitation* 2002;54:147–57.
21. Babbs CF, Nadkarni V. Optimizing chest compression to rescue ventilation ratios during onerescuer CPR by professionals and lay persons: children are not just little adults. *Resuscitation* 2004;61:173–81.
22. Kitamura T, Iwami T, Kawamura T, et al. Conventional and chestcompressiononly cardiopulmonary resuscitation by bystanders for children who have outofhospital cardiac arrests: a prospective, nationwide, populationbased cohort study. *Lancet* 2010.
23. Hourii PK, Frank LR, Menegazzi JJ, Taylor R. A randomized, controlled trial of twothumb vs twofinger chest compression in a swine infant model of cardiac arrest [see comment]. *Prehosp Emerg Care* 1997;1:65–7.
24. David R. Closed chest cardiac massage in the newborn infant. *Pediatrics* 1988;81:552–4.
25. Dorfsman ML, Menegazzi JJ, Wadas RJ, Auble TE. Twothumb vs twofinger chest compression in an infant model of prolonged cardiopulmonary resuscitation. *Acad Emerg Med* 2000;7:1077–82.
26. Whitelaw CC, Slywka B, Goldsmith LJ. Comparison of a twofinger versus twothumb method for chest compressions by healthcare providers in an infant mechanical model. *Resuscitation* 2000;43:213–6.
27. Menegazzi JJ, Auble TE, Nicklas KA, Hosack GM, Rack L, Goode JS. Twothumb versus twofinger chest compression during CRP in a swine infant model of cardiac arrest. *Ann Emerg Med* 1993;22:240–3.
28. Stevenson AG, McGowan J, Evans AL, Graham CA. CPR for children: one hand or two? *Resuscitation* 2005;64:205–8.
29. Gurnett CA, Atkins DL. Successful use of a biphasic waveform automated external defibrillator in a highrisk child. *Am J Cardiol* 2000;86:1051–3.
30. Konig B, Bengler J, Goldsworthy L. Automatic external defibrillation in a 6 year old. *Arch Dis Child* 2005;90:310–1.
31. Atkinson E, Mikysa B, Conway JA, et al. Specificity and sensitivity of automated external defibrillator rhythm analysis in infants and children. *Ann Emerg Med* 2003;42:185–96.
32. Cecchin F, Jorgenson DB, Berul CI, et al. Is arrhythmia detection by automatic external defibrillator accurate for children? Sensitivity and specificity of an automatic external defibrillator algorithm in 696 pediatric arrhythmias. *Circulation* 2001;103:2483–8.
33. Atkins DL, Scott WA, Blafox AD, et al. Sensitivity and specificity of an automated external defibrillator algorithm designed for pediatric patients. *Resuscitation* 2008;76:168–74.
34. Samson R, Berg R, Bingham R. Pediatric Advanced Life Support Task Force ILCOR. Use of automated external defibrillators for children: an update. An advisory statement from the Pediatric Advanced Life Support Task Force, International Liaison Committee on Resuscitation. *Resuscitation* 2003;57:237–43.
35. Jorgenson D, Morgan C, Snyder D, et al. Energy attenuator for pediatric application of an automated external defibrillator. *Crit Care Med* 2002;30: S145–7.
36. Tang W, Weil MH, Jorgenson D, et al. Fixedenergy biphasic waveform defibrillation in a pediatric model of cardiac arrest and resuscitation. *Crit Care Med* 2002;30:2736–41.
37. Berg RA, Chapman FW, Berg MD, et al. Attenuated adult biphasic shocks compared with weightbased monophasic shocks in a swine model of prolonged pediatric ventricular fibrillation. *Resuscitation* 2004;61:189–97.
38. Berg RA, Samson RA, BergMD, et al. Better outcome after pediatric defibrillation dosage than adult dosage in a swine model of pediatric ventricular fibrillation. *J Am Coll Cardiol* 2005;45:786–9.
39. BarCohen Y, Walsh EP, Love BA, Cecchin F. First appropriate use of automated external defibrillator in an infant. *Resuscitation* 2005;67:135–7.
40. Divekar A, Soni R. Successful parental use of an automated external defibrillator for an infant with longQT syndrome. *Pediatrics* 2006;118:e526–9.
41. Atkins DL, EversonStewart S, Sears GK, et al. Epidemiology and outcomes from outofhospital cardiac arrest in children: the Resuscitation Outcomes Consortium EpistryCardiac Arrest. *Circulation* 2009;119:1484–91.
42. RodriguezNunez A, LopezHerce J, Garcia C, Dominguez P, Carrillo A, Bellon JM. Pediatric defibrillation after cardiac arrest: initial response and outcome. *Crit Care* 2006;10:R113.
43. Samson RA, Nadkarni VM, Meaney PA, Carey SM, Berg MD, Berg RA. Outcomes of inhospital ventricular fibrillation in children. *N Engl J Med* 2006;354:2328–39.
44. Rea TD, Helbock M, Perry S, et al. Increasing use of cardiopulmonary resuscitation during outofhospital ventricular fibrillation arrest: survival implications of guideline changes. *Circulation* 2006;114:2760–5.
45. Menegazzi JJ, Hsieh M, Niemann JT, Swor RA. Derivation of clinical predictors of failed rescue shock during outofhospital ventricular fibrillation. *Prehosp Emerg Care* 2008;12:347–51.

46. Rea TD, Shah S, Kudenchuk PJ, Copass MK, Cobb LA. Automated external defibrillators: to what extent does the algorithm delay CPR? *Ann Emerg Med* 2005;46:132–41.
47. Becker L, Gold LS, Eisenberg M, White L, Hearne T, Rea T. Ventricular fibrillation in King County, Washington: a 30year perspective. *Resuscitation* 2008;79:22–7.
48. Rossano J, Quan L, Schiff M, MA K, DL A. Survival is not correlated with defibrillation dosing in pediatric outofhospital ventricular fibrillation. *Circulation* 2003;108. IV3201.
49. Clark CB, Zhang Y, Davies LR, Karlsson G, Kerber RE. Pediatric transthoracic defibrillation: biphasic versus monophasic waveforms in an experimental model. *Resuscitation* 2001;51:159–63.
50. Berg MD, Samson RA, Meyer RJ, Clark LL, Valenzuela TD, Berg RA. Pediatric defibrillation doses often fail to terminate prolonged outofhospital ventricular fibrillation in children. *Resuscitation* 2005;67:63–7.
51. Schneider T, Martens PR, Paschen H, et al. Multicenter, randomized, controlled trial of 150J biphasic shocks compared with 200to 360J monophasic shocks in the resuscitation of outofhospital cardiac arrest victims. Optimized Response to Cardiac Arrest (ORCA) Investigators. *Circulation* 2000;102:1780–7.
52. Faddy SC, Powell J, Craig JC. Biphasic and monophasic shocks for transthoracic defibrillation: a meta analysis of randomised controlled trials. *Resuscitation* 2003;58:9–16.
53. van Alem AP, Chapman FW, Lank P, Hart AA, Koster RW. A prospective, randomised and blinded comparison of first shock success of monophasic and biphasic waveforms in outofhospital cardiac arrest. *Resuscitation* 2003;58:17–24.
54. Safranek DJ, Eisenberg MS, Larsen MP. The epidemiology of cardiac arrest in young adults. *Ann Emerg Med* 1992;21:1102–6.
55. Redding JS. The choking controversy: critique of evidence on the Heimlich maneuver. *Crit Care Med* 1979;7:475–9.
56. Kuisma M, Suominen P, Korpela R. Paediatric outofhospital cardiac arrests – epidemiology and outcome. *Resuscitation* 1995;30:141–50.
57. Sirbaugh PE, Pepe PE, Shook JE, et al. A prospective, populationbased study of the demographics, epidemiology, management, and outcome of outofhospital pediatric cardiopulmonary arrest. *Ann Emerg Med* 1999;33:174–84.
58. Hickey RW, Cohen DM, Strausbaugh S, Dietrich AM. Pediatric patients requiring CPR in the prehospital setting. *Ann Emerg Med* 1995;25:495–501.
59. Young KD, Seidel JS. Pediatric cardiopulmonary resuscitation: a collective review. *Ann Emerg Med* 1999;33:195–205.
60. Reis AG, Nadkarni V, Perondi MB, Grisi S, Berg RA. A prospective investigation into the epidemiology of inhospital pediatric cardiopulmonary resuscitation using the international Utstein reporting style. *Pediatrics* 2002;109: 200–9.
61. Young KD, GauscheHill M, McClung CD, Lewis RJ. A prospective, populationbased study of the epidemiology and outcome of outofhospital pediatric cardiopulmonary arrest. *Pediatrics* 2004;114:157–64.
62. Richman PB, Nashed AH. The etiology of cardiac arrest in children and young adults: special considerations for ED management. *Am J Emerg Med* 1999;17:264–70.
63. Engdahl J, Bang A, Karlson BW, Lindqvist J, Herlitz J. Characteristics and outcome among patients suffering from out of hospital cardiac arrest of noncardiac aetiology. *Resuscitation* 2003;57:33–41.
64. Tibballs J, Kinney S. Reduction of hospital mortality and of preventable cardiac arrest and death on introduction of a pediatric medical emergency team. *Pediatr Crit Care Med* 2009;10:306–12.
65. Chan PS, Jain R, Nallmothu BK, Berg RA, Sasson C. Rapid response teams: a systematic review and metaanalysis. *Arch Intern Med* 2010;170:18–26.
66. Hunt EA, Zimmer KP, Rinke ML, et al. Transition from a traditional code team to a medical emergency team and categorization of cardiopulmonary arrests in a children’s center. *Arch Pediatr Adolesc Med* 2008;162:117–22.
67. Sharek PJ, Parast LM, Leong K, et al. Effect of a rapid response team on hospitalwide mortality and code rates outside the ICU in a Children’s Hospital. *JAMA* 2007;298:2267–74.
68. Brill R, Gibson R, Luria JW, et al. Implementation of a medical emergency team in a large pediatric teaching hospital prevents respiratory and cardiopulmonary arrests outside the intensive care unit. *Pediatr Crit Care Med* 2007;8:236–46, quiz 47.
69. Tibballs J, Kinney S, Duke T, Oakley E, Hennessy M. Reduction of paediatric inpatient cardiac arrest and death with a medical emergency team: preliminary results. *Arch Dis Child* 2005;90:1148–52.
70. Carcillo JA. Pediatric septic shock and multiple organ failure. *Crit Care Clin* 2003;19:413–40, viii.
71. Tibballs J, Russell P. Reliability of pulse palpation by healthcare personnel to diagnose paediatric cardiac arrest. *Resuscitation* 2009;80:61–4.

72. Eberle B, Dick WF, Schneider T, Wisser G, Doetsch S, Tzanova I. Checking the carotid pulse check: diagnostic accuracy of first responders in patients with and without a pulse. *Resuscitation* 1996;33:107–16.
73. Moule P. Checking the carotid pulse: diagnostic accuracy in students of the healthcare professions. *Resuscitation* 2000;44:195–201.
74. Lapostolle F, Le Toumelin P, Agostinucci JM, Catineau J, Adnet F. Basic cardiac life support providers checking the carotid pulse: performance, degree of conviction, and influencing factors. *Acad Emerg Med* 2004;11:878–80.
75. Frederick K, Bixby E, Orzel MN, StewartBrown S, Willett K. Will changing the emphasis from ‘pulseless’ to ‘no signs of circulation’ improve the recall scores for effective life support skills in children? *Resuscitation* 2002;55:255–61.
76. Tsung JW, Blaiwas M. Feasibility of correlating the pulse check with focused pointofcare echocardiography during pediatric cardiac arrest: a case series. *Resuscitation* 2008;77:264–9.
77. Dung NM, Day NPJ, Tam DTH, et al. Fluid replacement in dengue shock syndrome: a randomized, doubleblind comparison of four intravenousfluid regimens. *Clin Infect Dis* 1999;29:787–94.
78. Ngo NT, Cao XT, Kneen R, et al. Acute management of dengue shock syndrome: a randomized doubleblind comparison of 4 intravenous fluid regimens in the first hour. *Clin Infect Dis* 2001;32:204–13.
79. Wills BA, Nguyen MD, Ha TL, et al. Comparison of three fluid solutions for resuscitation in dengue shock syndrome. *N Engl J Med* 2005;353:877–89.
80. Upadhyay M, Singhi S, Murlidharan J, Kaur N, Majumdar S. Randomized evaluation of fluid resuscitation with crystalloid (saline) and colloid (polymer from degraded gelatin in saline) in pediatric septic shock. *Indian Pediatr* 2005;42:223–31.
81. Rechner JA, Loach VJ, Ali MT, Barber VS, Young JD, Mason DG. A comparison of the laryngeal mask airway with facemask and oropharyngeal airway for manual ventilation by critical care nurses in children. *Anaesthesia* 2007;62:790–5.
82. Blevin AE, McDouall SF, Rechner JA, et al. A comparison of the laryngeal mask airway with the facemask and oropharyngeal airway for manual ventilation by first responders in children. *Anaesthesia* 2009;64:1312–6.
83. Park C, Bahk JH, Ahn WS, Do SH, Lee KH. The laryngeal mask airway in infants and children. *Can J Anaesth* 2001;48:413–7.
84. Harnett M, Kinirons B, Heffernan A, Motherway C, Casey W. Airway complications in infants: comparison of laryngeal mask airway and the facemaskoral airway. *Can J Anaesth* 2000;47:315–8.
85. Scheller B, Schalk R, Byhahn C, et al. Laryngeal tube suction II for difficult airway management in neonates and small infants. *Resuscitation* 2009;80:805–10.
86. Hedges JR, Mann NC, Meischke H, Robbins M, Goldberg R, Zapka J. Assessment of chest pain onset and outofhospital delay using standardized interview questions: the REACT Pilot Study. Rapid Early Action for Coronary Treatment (REACT) Study Group. *Acad Emerg Med* 1998;5:773–80.
87. MurphyMacabobby M, Marshall WJ, Schneider C, Dries D. Neuromuscular blockade in aeromedical airway management. *AnnEmergMed* 1992;21:664–8.
88. Sayre M, Weisgerber I. The use of neuromuscular blocking agents by air medical services. *J Air Med Transp* 1992;11:7–11.
89. Rose W, Anderson L, Edmond S. Analysis of intubations. Before and after establishment of a rapid sequence intubation protocol for air medical use. *Air Med J* 1994;13:475–8.
90. Sing RF, Reilly PM, Rotondo MF, Lynch MJ, McCans JP, Schwab CW. Outofhospital rapidsequence induction for intubation of the pediatric patient. *Acad Emerg Med* 1996;3:41–5.
91. Ma OJ, Atchley RB, Hatley T, Green M, Young J, Brady W. Intubation success rates improve for an air medical program after implementing the use of neuromuscular blocking agents. *Am J Emerg Med* 1998;16:125–7.
92. Tayal V, Riggs R, Marx J, Tomaszewski C, Schneider R. Rapidsequence intubation at an emergency medicine residency: success rate and adverse events during a twoyear period. *Acad Emerg Med* 1999;6:31–7.
93. Wang HE, Sweeney TA, O’Connor RE, Rubinstein H. Failed prehospital intubations: an analysis of emergency department courses and outcomes. *Prehosp Emerg Care* 2001;5:134–41.
94. Kaye K, Frascone RJ, Held T. Prehospital rapidsequence intubation: a pilot training program. *Prehosp Emerg Care* 2003;7:235–40.
95. Wang HE, Kupas DF, Paris PM, Bates RR, Costantino JP, Yealy DM. Multivariate predictors of failed prehospital endotracheal intubation. *Acad Emerg Med* 2003;10:717–24.
96. Pepe P, Zachariah B, Chandra N. Invasive airway technique in resuscitation. *Ann Emerg Med* 1991;22:393–403.

97. Eich C, Roessler M, Nemeth M, Russo SG, Heuer JF, Timmermann A. Characteristics and outcome of prehospital paediatric tracheal intubation attended by anaesthesiatrained emergency physicians. *Resuscitation* 2009;80:1371–7.
98. Sagarin MJ, Chiang V, Sakles JC, et al. Rapid sequence intubation for pediatric emergency airway management. *Pediatr Emerg Care* 2002;18:417–23.
99. Moynihan RJ, BrockUtne JG, Archer JH, Feld LH, Kreitzman TR. The effect of cricoid pressure on preventing gastric insufflation in infants and children. *Anesthesiology* 1993;78:652–6.
100. Salem MR, Joseph NJ, Heyman HJ, Belani B, Paulissian R, Ferrara TP. Cricoid compression is effective in obliterating the esophageal lumen in the presence of a nasogastric tube. *Anesthesiology* 1985;63:443–6.
101. Walker RW, Ravi R, Haylett K. Effect of cricoid force on airway calibre in children: a bronchoscopic assessment. *Br J Anaesth* 2010;104:71–4.
102. Khine HH, Corddry DH, Kettrick RG, et al. Comparison of cuffed and uncuffed endotracheal tubes in young children during general anesthesia. *Anesthesiology* 1997;86:627–31, discussion 27A.
103. Weiss M, Dullenkopf A, Fischer JE, Keller C, Gerber AC. Prospective randomized controlled multicentre trial of cuffed or uncuffed endotracheal tubes in small children. *Br J Anaesth* 2009;103:867–73.
104. Duracher C, Schmutz E, Martinon C, Faivre J, Carli P, Orliaguet G. Evaluation of cuffed tracheal tube size predicted using the Khine formula in children. *Paediatr Anaesth* 2008;18:113–8.
105. Dullenkopf A, Kretschmar O, Knirsch W, et al. Comparison of tracheal tube cuff diameters with internal transverse diameters of the trachea in children. *Acta Anaesthesiol Scand* 2006;50:201–5.
106. Dullenkopf A, Gerber AC, Weiss M. Fit and seal characteristics of a new paediatric tracheal tube with high volumelow pressure polyurethane cuff. *Acta Anaesthesiol Scand* 2005;49:232–7.
107. Salgo B, Schmitz A, Henze G, et al. Evaluation of a new recommendation for improved cuffed tracheal tube size selection in infants and small children. *Acta Anaesthesiol Scand* 2006;50:557–61.
108. Luten RC, Wears RL, Broselow J, et al. Lengthbased endotracheal tube and emergency equipment in pediatrics. *Ann Emerg Med* 1992;21:900–4.
109. Deakers TW, Reynolds G, Stretton M, Newth CJ. Cuffed endotracheal tubes in pediatric intensive care. *J Pediatr* 1994;125:57–62.
110. Newth CJ, Rachman B, Patel N, Hammer J. The use of cuffed versus uncuffed endotracheal tubes in pediatric intensive care. *J Pediatr* 2004;144:333–7.
111. Dorsey DP, Bowman SM, Klein MB, Archer D, Sharar SR. Perioperative use of cuffed endotracheal tubes is advantageous in young pediatric burn patients. *Burns* 2010.
112. Mhanna MJ, Zamel YB, Tichy CM, Super DM. The “air leak” test around the endotracheal tube, as a predictor of postextubation stridor, is age dependent in children. *Crit Care Med* 2002;30:2639–43.
113. Katz SH, Falk JL. Misplaced endotracheal tubes by paramedics in an urban emergency medical services system. *Ann Emerg Med* 2001;37:32–7.
114. Gausche M, Lewis RJ, Stratton SJ, et al. Effect of outofhospital pediatric endotracheal intubation on survival and neurological outcome: a controlled clinical trial. *JAMA* 2000;283:783–90.
115. Kelly JJ, Eynon CA, Kaplan JL, de Garavilla L, Dalsey WC. Use of tube condensation as an indicator of endotracheal tube placement. *Ann Emerg Med* 1998;31:575–8.
116. Andersen KH, Hald A. Assessing the position of the tracheal tube: the reliability of different methods. *Anaesthesia* 1989;44:984–5.
117. Andersen KH, SchultzLebahn T. Oesophageal intubation can be undetected by auscultation of the chest. *Acta Anaesthesiol Scand* 1994;38:580–2.
118. Hartrey R, Kestin IG. Movement of oral and nasal tracheal tubes as a result of changes in head and neck position. *Anaesthesia* 1995;50:682–7.
119. Van de Louw A, Cracco C, Cerf C, et al. Accuracy of pulse oximetry in the intensive care unit. *Intensive Care Med* 2001;27:1606–13.
120. Seguin P, Le Rouzo A, Tanguy M, Guillou YM, Feuillu A, Malledant Y. Evidence for the need of bedside accuracy of pulse oximetry in an intensive care unit. *Crit Care Med* 2000;28:703–6.
121. Tan A, Schulze A, O’Donnell CP, Davis PG. Air versus oxygen for resuscitation of infants at birth. *Cochrane Database Syst Rev* 2004:CD002273.
122. Ramji S, Rasaily R, Mishra PK, et al. Resuscitation of asphyxiated newborns with room air or 100% oxygen at birth: a multicentric clinical trial. *Indian Pediatr* 2003;40:510–7.
123. Vento M, Asensi M, Sastre J, GarciaSala F, Pallardo FV, Vina J. Resuscitation with room air instead of 100% oxygen prevents oxidative stress in moderately asphyxiated term neonates. *Pediatrics* 2001;107:642–7.
124. Saugstad OD. Resuscitation of newborn infants with room air or oxygen. *Semin Neonatol* 2001;6:233–9.

125. Aufderheide TP, Sigurdsson G, Pirralo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation* 2004;109:1960–5.
126. Aufderheide TP, Lurie KG. Death by hyperventilation: a common and lifethreatening problem during cardiopulmonary resuscitation. *Crit Care Med* 2004;32:S345–51.
127. Wik L, Kramer-Johansen J, Myklebust H, et al. Quality of cardiopulmonary resuscitation during out-of-hospital cardiac arrest. *JAMA* 2005;293: 299–304.
128. Abella BS, Alvarado JP, Myklebust H, et al. Quality of cardiopulmonary resuscitation during in-hospital cardiac arrest. *JAMA* 2005;293:305–10.
129. Abella BS, Sandbo N, Vassilatos P, et al. Chest compression rates during cardiopulmonary resuscitation are suboptimal: a prospective study during in-hospital cardiac arrest. *Circulation* 2005;111:428–34.
130. Borke WB, Munkeby BH, Morkrid L, Thaulow E, Saugstad OD. Resuscitation with 100% O₂ does not protect the myocardium in hypoxic newborn piglets. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F156–60.
131. O’Neill JF, Deakin CD. Do we hyperventilate cardiac arrest patients? *Resuscitation* 2007;73:82–5.
132. Stockinger ZT, McSwain Jr NE. Prehospital endotracheal intubation for trauma does not improve survival over bag-valve-mask ventilation. *J Trauma* 2004;56:531–6.
133. Pitetti R, Glustein JZ, Bhende MS. Prehospital care and outcome of pediatric out-of-hospital cardiac arrest. *Prehosp Emerg Care* 2002;6:283–90.
134. Cooper A, DiScala C, Foltin G, Tunik M, Markenson D, Welborn C. Prehospital endotracheal intubation for severe head injury in children: a reappraisal. *Semin Pediatr Surg* 2001;10:3–6.
135. DiRusso SM, Sullivan T, Risucci D, Nealon P, Slim M. Intubation of pediatric trauma patients in the field: predictor of negative outcome despite risk stratification. *J Trauma* 2005;59:84–90, discussion – 1.
136. Bhende MS, Thompson AE, Orr RA. Utility of an endtidal carbon dioxide detector during stabilization and transport of critically ill children. *Pediatrics* 1992;89:1042–4.
137. Bhende MS, LaCovey DC. Endtidal carbon dioxide monitoring in the prehospital setting. *Prehosp Emerg Care* 2001;5:208–13.
138. Ornato JP, Shipley JB, Racht EM, et al. Multicenter study of a portable, handsize, colorimetric endtidal carbon dioxide detection device. *Ann Emerg Med* 1992;21:518–23.
139. Gonzalez del Rey JA, Poirier MP, Digiulio GA. Evaluation of an ambubag valve with a self-contained, colorimetric endtidal CO₂ system in the detection of airway mishaps: an animal trial. *Pediatr Emerg Care* 2000;16:121–3.
140. Bhende MS, Thompson AE. Evaluation of an endtidal CO₂ detector during pediatric cardiopulmonary resuscitation. *Pediatrics* 1995;95:395–9.
141. Bhende MS, Karasic DG, Karasic RB. Endtidal carbon dioxide changes during cardiopulmonary resuscitation after experimental asphyxial cardiac arrest. *Am J Emerg Med* 1996;14:349–50.
142. DeBehnke DJ, Hilander SJ, Dobler DW, Wickman LL, Swart GL. The hemodynamic and arterial blood gas response to asphyxiation: a canine model of pulseless electrical activity. *Resuscitation* 1995;30:169–75.
143. Ornato JP, Garnett AR, Glauser FL. Relationship between cardiac output and the endtidal carbon dioxide tension. *Ann Emerg Med* 1990;19:1104–6.
144. Mauer D, Schneider T, Elich D, Dick W. Carbon dioxide levels during prehospital active compression – decompression versus standard cardiopulmonary resuscitation. *Resuscitation* 1998;39:67–74.
145. Kolar M, Krizmaric M, Klemen P, Grmec S. Partial pressure of endtidal carbon dioxide successfully predicts cardiopulmonary resuscitation in the field: a prospective observational study. *Crit Care* 2008;12:R115.
146. Callahan M, Barton C, Matthay M. Effect of epinephrine on the ability of endtidal carbon dioxide readings to predict initial resuscitation from cardiac arrest. *Crit Care Med* 1992;20:337–43.
147. Cantineau JP, Merckx P, Lambert Y, Sorkine M, Bertrand C, Duvaldestin P. Effect of epinephrine on endtidal carbon dioxide pressure during prehospital cardiopulmonary resuscitation. *Am J Emerg Med* 1994;12:267–70.
148. Chase PB, Kern KB, Sanders AB, Otto CW, Ewy GA. Effects of graded doses of epinephrine on both noninvasive and invasive measures of myocardial perfusion and blood flow during cardiopulmonary resuscitation. *Crit Care Med* 1993;21:413–9.
149. Gonzalez ER, Ornato JP, Garnett AR, Levine RL, Young DS, Racht EM. Dose-dependent vasopressor response to epinephrine during CPR in human beings. *Ann Emerg Med* 1989;18:920–6.
150. Lindberg L, Liao Q, Steen S. The effects of epinephrine/norepinephrine on endtidal carbon dioxide concentration, coronary perfusion pressure and pulmonary arterial blood flow during cardiopulmonary resuscitation. *Resuscitation* 2000;43:129–40.

151. Falk JL, Rackow EC, Weil MH. Endtidal carbon dioxide concentration during cardiopulmonary resuscitation. *N Engl J Med* 1988;318:607–11.
152. Sharieff GQ, Rodarte A, Wilton N, Bleyle D. The selfinflating bulb as an airway adjunct: is it reliable in children weighing less than 20 kilograms? *Acad Emerg Med* 2003;10:303–8.
153. Sharieff GQ, Rodarte A, Wilton N, Silva PD, Bleyle D. The selfinflating bulb as an esophageal detector device in children weighing more than twenty kilograms: a comparison of two techniques. *Ann Emerg Med* 2003;41:623–9.
154. Poirier MP, Gonzalez DelRey JA, McAneney CM, DiGiulio GA. Utility of monitoring capnography, pulse oximetry, and vital signs in the detection of airway mishaps: a hyperoxemic animal model. *Am J Emerg Med* 1998;16:350–2.
155. Lillis KA, Jaffe DM. Prehospital intravenous access in children. *Ann Emerg Med* 1992;21:1430–4.
156. Neufeld JD, Marx JA, Moore EE, Light AI. Comparison of intraosseous, central, and peripheral routes of crystalloid infusion for resuscitation of hemorrhagic shock in a swine model. *J Trauma* 1993;34:422–8.
157. Hedges JR, Barsan WB, Doan LA, et al. Central versus peripheral intravenous routes in cardiopulmonary resuscitation. *Am J Emerg Med* 1984;2:385–90.
158. Kanter RK, Zimmerman JJ, Strauss RH, Stoeckel KA. Pediatric emergency intravenous access. Evaluation of a protocol. *Am J Dis Child* 1986;140:132–4.
159. Banerjee S, Singhi SC, Singh S, Singh M. The intraosseous route is a suitable alternative to intravenous route for fluid resuscitation in severely dehydrated children. *Indian Pediatr* 1994;31:1511–20.
160. Glaeser PW, Hellmich TR, Szewczuga D, Losek JD, Smith DS. Fiveyear experience in prehospital intraosseous infusions in children and adults. *Ann Emerg Med* 1993;22:1119–24.
161. Guy J, Haley K, Zuspan SJ. Use of intraosseous infusion in the pediatric trauma patient. *J Pediatr Surg* 1993;28:158–61.
162. Orłowski JP, Julius CJ, Petras RE, Porembka DT, Gallagher JM. The safety of intraosseous infusions: risks of fat and bone marrow emboli to the lungs. *Ann Emerg Med* 1989;18:1062–7.
163. Orłowski JP, Porembka DT, Gallagher JM, Lockrem JD, VanLente F. Comparison study of intraosseous, central intravenous, and peripheral intravenous infusions of emergency drugs. *Am J Dis Child* 1990;144:112–7.
164. Abe KK, Blum GT, Yamamoto LG. Intraosseous is faster and easier than umbilical venous catheterization in newborn emergency vascular access models. *Am J Emerg Med* 2000;18:126–9.
165. Ellemunter H, Simma B, Trawoger R, Maurer H. Intraosseous lines in preterm and full term neonates. *Arch Dis Child Fetal Neonatal Ed* 1999;80:F74–5.
166. Fiorito BA, Mirza F, Doran TM, et al. Intraosseous access in the setting of pediatric critical care transport. *Pediatr Crit Care Med* 2005;6:50–3.
167. Horton MA, Beamer C. Powered intraosseous insertion provides safe and effective vascular access for pediatric emergency patients. *Pediatr Emerg Care* 2008;24:347–50.
168. Frascone RJ, Jensen J, Wewerka SS, Salzman JG. Use of the pediatric EZIO needle by emergency medical services providers. *Pediatr Emerg Care* 2009;25: 329–32.
169. Cameron JL, Fontanarosa PB, Passalacqua AM. A comparative study of peripheral to central circulation delivery times between intraosseous and intravenous injection using a radionuclide technique in normovolemic and hypovolemic canines. *J Emerg Med* 1989;7:123–7.
170. Warren DW, Kisson N, Sommerauer JF, Rieder MJ. Comparison of fluid infusion rates among peripheral intravenous and humerus, femur, malleolus, and tibial intraosseous sites in normovolemic and hypovolemic piglets. *Ann Emerg Med* 1993;22:183–6.
171. Brickman KR, Krupp K, Rega P, Alexander J, Guinness M. Typing and screening of blood from intraosseous access. *Ann Emerg Med* 1992;21:414–7.
172. Johnson L, Kisson N, Fiallos M, Abdelmoneim T, Murphy S. Use of intraosseous blood to assess blood chemistries and hemoglobin during cardiopulmonary resuscitation with drug infusions. *Crit Care Med* 1999;27:1147–52.
173. Ummenhofer W, Frei FJ, Urwyler A, Drewe J. Are laboratory values in bone marrow aspirate predictable for venous blood in paediatric patients? *Resuscitation* 1994;27:123–8.
174. Abdelmoneim T, Kisson N, Johnson L, Fiallos M, Murphy S. Acidbase status of blood from intraosseous and mixed venous sites during prolonged cardiopulmonary resuscitation and drug infusions. *Crit Care Med* 1999;27:1923–8.
175. Voelckel WG, Lindner KH, Wenzel V, et al. Intraosseous blood gases during hypothermia: correlation with arterial, mixed venous, and sagittal sinus blood. *Crit Care Med* 2000;28:2915–20.

176. Kissoon N, Peterson R, Murphy S, Gayle M, Ceithaml E, HarwoodNuss A. Comparison of pH and carbon dioxide tension values of central venous and intraosseous blood during changes in cardiac output. *Crit Care Med* 1994;22:1010–5.
177. Eisenkraft A, Gilat E, Chapman S, Baranes S, Egoz I, Levy A. Efficacy of the bone injection gun in the treatment of organophosphate poisoning. *Biopharm Drug Dispos* 2007;28:145–50.
178. Brenner T, Bernhard M, Helm M, et al. Comparison of two intraosseous infusion systems for adult emergency medical use. *Resuscitation* 2008;78:314–9.
179. Venkataraman ST, Orr RA, Thompson AE. Percutaneous infraclavicular subclavian vein catheterization in critically ill infants and children. *J Pediatr* 1988;113:480–5.
180. Fleisher G, Caputo G, Baskin M. Comparison of external jugular and peripheral venous administration of sodium bicarbonate in puppies. *Crit Care Med* 1989;17:251–4.
181. Stenzel JP, Green TP, Fuhrman BP, Carlson PE, Marchessault RP. Percutaneous femoral venous catheterizations: a prospective study of complications. *J Pediatr* 1989;114:411–5.
182. Kleinman ME, OhW, Stonestreet BS. Comparison of intravenous and endotracheal epinephrine during cardiopulmonary resuscitation in newborn piglets. *Crit Care Med* 1999;27:2748–54.
183. Hahnel JH, Lindner KH, Schurmann C, Prengel A, Ahnefeld FW. Plasma lidocaine levels and PaO₂ with endobronchial administration: dilution with normal saline or distilled water? *Ann Emerg Med* 1990;19:1314–7.
184. Jasani MS, Nadkarni VM, Finkelstein MS, Mandell GA, Salzman SK, Norman ME. Effects of different techniques of endotracheal epinephrine administration in pediatric porcine hypoxichypercarbic cardiopulmonary arrest. *Crit Care Med* 1994;22:1174–80.
185. Steinfath M, Scholz J, Schulte am Esch J, Laer S, Reymann A, Scholz H. The technique of endobronchial lidocaine administration does not influence plasma concentration profiles and pharmacokinetic parameters in humans. *Resuscitation* 1995;29:55–62.
186. Carcillo JA, Fields AI. Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med* 2002;30:1365–78.
187. Simma B, Burger R, Falk M, Sacher P, Fanconi S. A prospective, randomized, and controlled study of fluid management in children with severe head injury: lactated Ringer's solution versus hypertonic saline. *Crit Care Med* 1998;26:1265–70.
188. Myburgh J, Cooper DJ, Finfer S, et al. Saline or albumin for fluid resuscitation in patients with traumatic brain injury. *N Engl J Med* 2007;357:874–84.
189. Rocha E, Silva M. Hypertonic saline resuscitation. *Medicina* 1998;58: 393–402.
190. Katz LM, Wang Y, Ebmeyer U, Radovsky A, Safar P. Glucose plus insulin infusion improves cerebral outcome after asphyxial cardiac arrest. *Neuroreport* 1998;9:3363–7.
191. Longstreth Jr WT, Copass MK, Dennis LK, RauchMatthews ME, Stark MS, Cobb LA. Intravenous glucose after outofhospital cardiopulmonary arrest: a communitybased randomized trial. *Neurology* 1993;43:2534–41.
192. Chang YS, Park WS, Ko SY, et al. Effects of fasting and insulininduced hypoglycemia on brain cell membrane function and energy metabolism during hypoxiaischemia in newborn piglets. *Brain Res* 1999;844:135–42.
193. Cherian L, Goodman JC, Robertson CS. Hyperglycemia increases brain injury caused by secondary ischemia after cortical impact injury in rats. *Crit Care Med* 1997;25:1378–83.
194. Paul T, Bertram H, Bokenkamp R, Hausdorf G. Supraventricular tachycardia in infants, children and adolescents: diagnosis, and pharmacological and interventional therapy. *Paediatr Drugs* 2000;2:171–81.
195. Losek JD, Endom E, Dietrich A, Stewart G, Zempsky W, Smith K. Adenosine and pediatric supraventricular tachycardia in the emergency department: multicenter study and review. *Ann Emerg Med* 1999;33:185–91.
196. Roberts JR, Greenburg MI, Knaub M, Baskin SI. Comparison of the pharmacological effects of epinephrine administered by the intravenous and endotracheal routes. *JACEP* 1978;7:260–4.
197. Zaritsky A. Pediatric resuscitation pharmacology. Members of the Medications in Pediatric Resuscitation Panel. *Ann Emerg Med* 1993;22:445–55.
198. Manisterski Y, Vaknin Z, BenAbraham R, et al. Endotracheal epinephrine: a call for larger doses. *Anesth Analg* 2002;95:1037–41, table of contents.
199. Efrati O, BenAbraham R, Barak A, et al. Endobronchial adrenaline: should it be reconsidered? Dose response and haemodynamic effect in dogs. *Resuscitation* 2003;59:117–22.
200. PattersonMD, Boenning DA, Klein BL, et al. The use of highdose epinephrine for patients with outofhospital cardiopulmonary arrest refractory to prehospital interventions. *Pediatr Emerg Care* 2005;21:227–37.

201. Perondi MB, Reis AG, Paiva EF, Nadkarni VM, Berg RA. A comparison of highdose and standarddose epinephrine in children with cardiac arrest. *N Engl J Med* 2004;350:1722–30.
202. Carpenter TC, Stenmark KR. Highdose epinephrine is not superior to standarddose epinephrine in pediatric in-hospital cardiopulmonary arrest. *Pediatrics* 1997;99:403–8.
203. Dieckmann RA, Vardis R. Highdose epinephrine in pediatric out-of-hospital cardiopulmonary arrest. *Pediatrics* 1995;95:901–13.
204. Berg RA, Otto CW, Kern KB, et al. Highdose epinephrine results in greater early mortality after resuscitation from prolonged cardiac arrest in pigs: a prospective, randomized study. *Crit Care Med* 1994;22:282–90.
205. Rubertsson S, Wiklund L. Hemodynamic effects of epinephrine in combination with different alkaline buffers during experimental, openchest, cardiopulmonary resuscitation. *Crit Care Med* 1993;21:1051–7.
206. Somberg JC, Timar S, Bailin SJ, et al. Lack of a hypotensive effect with rapid administration of a new aqueous formulation of intravenous amiodarone. *Am J Cardiol* 2004;93:576–81.
207. Yap SC, Hoomtje T, Sreeram N. Polymorphic ventricular tachycardia after use of intravenous amiodarone for postoperative junctional ectopic tachycardia. *Int J Cardiol* 2000;76:245–7.
208. Dauchot P, Gravenstein JS. Effects of atropine on the electrocardiogram in different age groups. *Clin Pharmacol Ther* 1971;12:274–80.
209. Yilmaz O, Eser M, Sahiner A, Altintop L, Yesildag O. Hypotension, bradycardia and syncope caused by honey poisoning. *Resuscitation* 2006;68:405–8.
210. Brady WJ, Swart G, DeBehnke DJ, Ma OJ, Aufderheide TP. The efficacy of atropine in the treatment of hemodynamically unstable bradycardia and atrioventricular block: prehospital and emergency department considerations. *Resuscitation* 1999;41:47–55.
211. Smith I, Monk TG, White PF. Comparison of transesophageal atrial pacing with anticholinergic drugs for the treatment of intraoperative bradycardia. *Anesth Analg* 1994;78:245–52.
212. Chadda KD, Lichstein E, Gupta PK, Kourtesis P. Effects of atropine in patients with bradyarrhythmia complicating myocardial infarction: usefulness of an optimum dose for overdrive. *Am J Med* 1977;63:503–10.
213. Stulz PM, Scheidegger D, Drop LJ, Lowenstein E, Laver MB. Ventricular pump performance during hypocalcemia: clinical and experimental studies. *J Thorac Cardiovasc Surg* 1979;78:185–94.
214. van Walraven C, Stiell IG, Wells GA, Hebert PC, Vandemheen K. Do advanced cardiac life support drugs increase resuscitation rates from in-hospital cardiac arrest? The OTAC Study Group. *Ann Emerg Med* 1998;32:544–53.
215. Paraskos JA. Cardiovascular pharmacology III: atropine, calcium, calcium blockers, and (beta)blockers. *Circulation* 1986;74:IV86–9.
216. Stueven HA, Thompson B, Aprahamian C, Tonsfeldt DJ, Kastenson EH. The effectiveness of calcium chloride in refractory electromechanical dissociation. *Ann Emerg Med* 1985;14:626–9.
217. Stueven HA, Thompson B, Aprahamian C, Tonsfeldt DJ, Kastenson EH. Lack of effectiveness of calcium chloride in refractory asystole. *Ann Emerg Med* 1985;14:630–2.
218. Srinivasan V, Morris MC, Helfaer MA, Berg RA, Nadkarni VM. Calcium use during in-hospital pediatric cardiopulmonary resuscitation: a report from the National Registry of Cardiopulmonary Resuscitation. *Pediatrics* 2008;121:e1144–51.
219. de Mos N, van Litsenburg RR, McCrindle B, Bohn DJ, Parshuram CS. Pediatric intensive care unit cardiac arrest: incidence, survival, and predictive factors. *Crit Care Med* 2006;34:1209–15.
220. Meert KL, Donaldson A, Nadkarni V, et al. Multicenter cohort study of in-hospital pediatric cardiac arrest. *Pediatr Crit Care Med* 2009;10:544–53.
221. Srinivasan V, Spinella PC, Drott HR, Roth CL, Helfaer MA, Nadkarni V. Association of timing, duration, and intensity of hyperglycemia with intensive care unit mortality in critically ill children. *Pediatr Crit Care Med* 2004;5:329–36.
222. Krinsley JS. Effect of an intensive glucose management protocol on the mortality of critically ill adult patients. *Mayo Clin Proc* 2004;79:992–1000.
223. Losek JD. Hypoglycemia and the ABC'S (sugar) of pediatric resuscitation. *Ann Emerg Med* 2000;35:43–6.
224. Finney SJ, Zekveld C, Elia A, Evans TW. Glucose control and mortality in critically ill patients. *JAMA* 2003;290:2041–7.
225. Losert H, Sterz F, Roine RO, et al. Strict normoglycaemic blood glucose levels in the therapeutic management of patients within 12 h after cardiac arrest might not be necessary. *Resuscitation* 2008;76:214–20.

226. Oksanen T, Skrifvars MB, Varpula T, et al. Strict versus moderate glucose control after resuscitation from ventricular fibrillation. *Intensive Care Med* 2007;33:2093–100.
227. Beardsall K, Vanhaesebrouck S, OgilvyStuart AL, et al. Early insulin therapy in verylowbirthweight infants. *N Engl J Med* 2008;359:1873–84.
228. Vlasselaers D, Milants I, Desmet L, et al. Intensive insulin therapy for patients in paediatric intensive care: a prospective, randomised controlled study. *Lancet* 2009;373:547–56.
229. Gandhi GY, Murad MH, Flynn DN, et al. Effect of perioperative insulin infusion on surgical morbidity and mortality: systematic review and metaanalysis of randomized trials. *Mayo Clin Proc* 2008;83:418–30.
230. Griesdale DE, de Souza RJ, van Dam RM, et al. Intensive insulin therapy and mortality among critically ill patients: a metaanalysis including NICESUGAR study data. *CMAJ* 2009;180:821–7.
231. Finfer S, Chittock DR, Su SY, et al. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;360:1283–97.
232. Allegra J, Lavery R, Cody R, et al. Magnesium sulfate in the treatment of refractory ventricular fibrillation in the prehospital setting. *Resuscitation* 2001;49:245–9.
233. Tzivoni D, Banai S, Schuger C, et al. Treatment of torsade de pointes with magnesium sulfate. *Circulation* 1988;77:392–7.
234. Lokesh L, Kumar P, Murki S, Narang A. A randomized controlled trial of sodium bicarbonate in neonatal resuscitationeffect on immediate outcome. *Resuscitation* 2004;60:219–23.
235. BarJoseph G, Abramson NS, Kelsey SF, Mashiach T, Craig MT, Safar P. Improved resuscitation outcome in emergency medical systems with increased usage of sodium bicarbonate during cardiopulmonary resuscitation. *Acta Anaesthesiol Scand* 2005;49:6–15.
236. Dorian P, Cass D, Schwartz B, Cooper R, Gelaznikas R, Barr A. Amiodarone as compared with lidocaine for shockresistant ventricular fibrillation. *N Engl J Med* 2002;346:884–90.
237. Walsh EP, Saul JP, Sholler GF, et al. Evaluation of a staged treatment protocol for rapid automatic junctional tachycardia after operation for congenital heart disease. *J Am Coll Cardiol* 1997;29:1046–53.
238. WangJD, Fu YC, Jan SL, Chi CS. Verapamil sensitive idiopathic ventricular tachycardia in an infant. *Jpn Heart J* 2003;44:667–71.
239. Chang PM, Silka MJ, Moromisato DY, BarCohen Y. Amiodarone versus procainamide for the acute treatment of recurrent supraventricular tachycardia in pediatric patients. *Circ Arrhythm Electrophysiol* 2010;3:134–40.
240. Singh BN, Kehoe R, Woosley RL, Scheinman M, Quart B. Multicenter trial of sotalol compared with procainamide in the suppression of inducible ventricular tachycardia: a doubleblind, randomized parallel evaluation. Sotalol Multicenter Study Group. *Am Heart J* 1995;129:87–97.
241. Luedtke SA, Kuhn RJ, McCaffrey FM. Pharmacologic management of supraventricular tachycardias in children. Part 1: WolffParkinsonWhite and atrioventricular nodal reentry. *Ann Pharmacother* 1997;31:1227–43.
242. Luedtke SA, Kuhn RJ, McCaffrey FM. Pharmacologic management of supraventricular tachycardias in children. Part 2: Atrial flutter, atrial fibrillation, and junctional and atrial ectopic tachycardia. *Ann Pharmacother* 1997;31: 1347–59.
243. Mandapati R, Byrum CJ, Kavey RE, et al. Procainamide for rate control of postsurgical junctional tachycardia. *Pediatr Cardiol* 2000;21:123–8.
244. Wang JN,WuJM, Tsai YC, Lin CS. Ectopic atrial tachycardia in children. *J Formos Med Assoc* 2000;99:766–70.
245. Wang R, Schuyler J, Raymond R. The role of the cell membrane bicarbonate exchanger in NaHCO₃ therapy of imipramine cardiac dysfunction. *J Toxicol Clin Toxicol* 1997;35:533.
246. Holmes CL, Landry DW, Granton JT. Science review: vasopressin and the cardiovascular system. Part 1 – Receptor physiology. *Crit Care* 2003;7: 427–34.
247. Voelckel WG, Lurie KG, McKnite S, et al. Effects of epinephrine and vasopressin in a piglet model of prolonged ventricular fibrillation and cardiopulmonary resuscitation. *Crit Care Med* 2002;30:957–62.
248. Voelckel WG, Lurie KG, McKnite S, et al. Comparison of epinephrine and vasopressin in a pediatric porcine model of asphyxial cardiac arrest. *Crit Care Med* 2000;28:3777–83.
249. Mann K, Berg RA, Nadkarni V. Beneficial effects of vasopressin in prolonged pediatric cardiac arrest: a case series. *Resuscitation* 2002;52:149–56.
250. Duncan JM, Meaney P, Simpson P, Berg RA, Nadkarni V, Schexnayder S. Vasopressin for inhospital pediatric cardiac arrest: results from the American Heart Association National Registry of Cardiopulmonary Resuscitation. *Pediatr Crit Care Med* 2009;10:191–5.
251. Callaway CW, Hostler D, Doshi AA, et al. Usefulness of vasopressin administered with epinephrine during outofhospital cardiac arrest. *Am J Cardiol* 2006;98:1316–21.

252. Gueugniaud PY, David JS, Chanzy E, et al. Vasopressin and epinephrine vs. epinephrine alone in cardiopulmonary resuscitation. *N Engl J Med* 2008;359:21–30.
253. Mukoyama T, Kinoshita K, Nagao K, Tanjoh K. Reduced effectiveness of vasopressin in repeated doses for patients undergoing prolonged cardiopulmonary resuscitation. *Resuscitation* 2009;80:755–61.
254. Aung K, Htay T. Vasopressin for cardiac arrest: a systematic review and metaanalysis. *Arch Intern Med* 2005;165:17–24.
255. Matok I, Vard A, Efrati O, et al. Terlipressin as rescue therapy for intractable hypotension due to septic shock in children. *Shock* 2005;23: 305–10.
256. Peters MJ, Booth RA, Petros AJ. Terlipressin bolus induces systemic vasoconstriction in septic shock. *Pediatr Crit Care Med* 2004;5:112–5.
257. RodriguezNunez A, LopezHerce J, GilAnton J, Hernandez A, Rey C. Rescue treatment with terlipressin in children with refractory septic shock: a clinical study. *Crit Care* 2006;10:R20.
258. GilAnton J, LopezHerce J, Morteruel E, Carrillo A, RodriguezNunez A. Pediatric cardiac arrest refractory to advanced life support: is there a role for terlipressin? *Pediatr Crit Care Med* 2010;11:139–41.
259. RodriguezNunez A, FernandezSanmartin M, MartinonTorres F, GonzalezAlonso N, MartinonSanchez JM. Terlipressin for catecholamineresistant septic shock in children. *Intensive Care Med* 2004;30:477–80.
260. Yildizdas D, Yapicioglu H, Celik U, Sertdemir Y, Alhan E. Terlipressin as a rescue therapy for catecholamineresistant septic shock in children. *Intensive Care Med* 2008;34:511–7.
261. Matok I, Vardi A, Augarten A, et al. Beneficial effects of terlipressin in prolonged pediatric cardiopulmonary resuscitation: a case series. *Crit Care Med* 2007;35:1161–4.
262. Atkins DL, Kerber RE. Pediatric defibrillation: current flow is improved by using “adult” electrode paddles. *Pediatrics* 1994;94:90–3.
263. Atkins DL, Sirna S, Kieso R, Charbonnier F, Kerber RE. Pediatric defibrillation: importance of paddle size in determining transthoracic impedance. *Pediatrics* 1988;82:914–8.
264. Deakin C, Sado D, Petley G, Clewlow F. Determining the optimal paddle force for external defibrillation. *Am J Cardiol* 2002;90:812–3.
265. Bennetts SH, Deakin CD, Petley GW, Clewlow F. Is optimal paddle force applied during paediatric external defibrillation? *Resuscitation* 2004;60:29–32.
266. Berg MD, Banville IL, Chapman FW, et al. Attenuating the defibrillation dosage decreases postresuscitation myocardial dysfunction in a swine model of pediatric ventricular fibrillation. *Pediatr Crit Care Med* 2008;9:429–34.
267. Atkins DL, Hartley LL, York DK. Accurate recognition and effective treatment of ventricular fibrillation by automated external defibrillators in adolescents. *Pediatrics* 1998;101:393–7.
268. Edelson DP, Abella BS, KramerJohansen J, et al. Effects of compression depth and preshock pauses predict defibrillation failure during cardiac arrest. *Resuscitation* 2006;71:137–45.
269. Eftestol T, Sunde K, Steen PA. Effects of interrupting precordial compressions on the calculated probability of defibrillation success during outofhospital cardiac arrest. *Circulation* 2002;105:2270–3.
270. Saul JP, Scott WA, Brown S, et al. Intravenous amiodarone for incessant tachyarrhythmias in children: a randomized, doubleblind, antiarrhythmic drug trial. *Circulation* 2005;112:3470–7.
271. Pierpont GL, Kruse JA, Nelson DH. Intraarterial monitoring during cardiopulmonary resuscitation. *Catheterization Cardiovasc Diagn* 1985;11: 513–20.
272. Steiger HV, Rimbach K, Muller E, Breikreutz R. Focused emergency echocardiography: lifesaving tool for a 14yearold girl suffering outofhospital pulseless electrical activity arrest because of cardiac tamponade. *Eur J Emerg Med* 2009;16:103–5.
273. Zaritsky A, Nadkarni V, Getson P, Kuehl K. CPR in children. *Ann Emerg Med* 1987;16:1107–11.
274. Mogayzel C, Quan L, Graves JR, Tiedeman D, Fahrenbruch C, Herndon P. Outofhospital ventricular fibrillation in children and adolescents: causes and outcomes. *Ann Emerg Med* 1995;25:484–91.
275. Herlitz J, Engdahl J, Svensson L, Young M, Angquist KA, Holmberg S. Characteristics and outcome among children suffering from out of hospital cardiac arrest in Sweden. *Resuscitation* 2005;64:37–40.
276. Berg RA. Role of mouthtomouth rescue breathing in bystander cardiopulmonary resuscitation for asphyxial cardiac arrest. *Crit Care Med* 2000;28(Suppl.):N193–5.
277. Kuisma M, Suominen P, Korpela R. Paediatric outofhospital cardiac arrests: epidemiology and outcome. *Resuscitation* 1995;30:141–50.
278. Kyriacou DN, Arcinue EL, Peek C, Kraus JF. Effect of immediate resuscitation on children with submersion injury. *Pediatrics* 1994;94:137–42.
279. Berg RA, Hilwig RW, Kern KB, Ewy GA. Bystander” chest compressions and assisted ventilation independently improve outcome from piglet asphyxial pulseless “cardiac arrest”. *Circulation* 2000;101:1743–8.

280. Appleton GO, Cummins RO, Larson MP, Graves JR. CPR and the single rescuer: at what age should you "call first" rather than "call fast"? *Ann Emerg Med* 1995;25:492-4.
281. Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predicting survival from outofhospital cardiac arrest: a graphic model. *Ann Emerg Med* 1993;22:1652-8.
282. Cobb LA, Fahrenbruch CE, Walsh TR, et al. Influence of cardiopulmonary resuscitation prior to defibrillation in patients with outofhospital ventricular fibrillation. *JAMA* 1999;281:1182-8.
283. Wik L, Hansen TB, Fylling F, et al. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with outofhospital ventricular fibrillation: a randomized trial. *JAMA* 2003;289:1389-95.
284. Jacobs IG, Finn JC, Ozer HF, Jelinek GA. CPR before defibrillation in outofhospital cardiac arrest: a randomized trial. *Emerg Med Australas* 2005;17:39-45.
285. Somberg JC, Bailin SJ, Haffajee CI, et al. Intravenous lidocaine versus intravenous amiodarone (in a new aqueous formulation) for incessant ventricular tachycardia. *Am J Cardiol* 2002;90:853-9.
286. Kudenchuk PJ, Cobb LA, Copass MK, et al. Amiodarone for resuscitation after outofhospital cardiac arrest due to ventricular fibrillation. *N Engl J Med* 1999;341:871-8.
287. Perry JC, Fenrich AL, Hulse JE, Triedman JK, Friedman RA, Lamberti JJ. Pediatric use of intravenous amiodarone: efficacy and safety in critically ill patients from a multicenter protocol. *J Am Coll Cardiol* 1996;27:1246-50.
288. Cummins RO, Graves JR, Larsen MP, et al. Outofhospital transcutaneous pacing by emergency medical technicians in patients with asystolic cardiac arrest. *N Engl J Med* 1993;328:1377-82.
289. Sreeram N, Wren C. Supraventricular tachycardia in infants: response to initial treatment. *Arch Dis Child* 1990;65:127-9.
290. Bianconi L, Castro A, Dinelli M, et al. Comparison of intravenously administered dofetilide versus amiodarone in the acute termination of atrial fibrillation and flutter. A multicentre, randomized, doubleblind, placebocontrolled study. *Eur Heart J* 2000;21:1265-73.
291. Burri S, Hug MI, Bauersfeld U. Efficacy and safety of intravenous amiodarone for incessant tachycardias in infants. *Eur J Pediatr* 2003;162:880-4.
292. Celiker A, Ceviz N, Ozme S. Effectiveness and safety of intravenous amiodarone in drugresistant tachyarrhythmias of children. *Acta Paediatr Jpn* 1998;40:567-72.
293. DodgeKhatami A, Miller O, Anderson R, GilJaurena J, Goldman A, de Leval M. Impact of junctional ectopic tachycardia on postoperative morbidity following repair of congenital heart defects. *Eur J Cardiothorac Surg* 2002;21:255-9.
294. Figa FH, Gow RM, Hamilton RM, Freedom RM. Clinical efficacy and safety of intravenous Amiodarone in infants and children. *Am J Cardiol* 1994;74:573-7.
295. Hoffman TM, Bush DM, Wernovsky G, et al. Postoperative junctional ectopic tachycardia in children: incidence, risk factors, and treatment. *Ann Thorac Surg* 2002;74:1607-11.
296. Soult JA, MunozM, Lopez JD, Romero A, Santos J, Tovaruela A. Efficacy and safety of intravenous amiodarone for shortterm treatment of paroxysmal supraventricular tachycardia in children. *Pediatr Cardiol* 1995;16:16-9.
297. Haas NA, Camphausen CK. Acute hemodynamic effects of intravenous amiodarone treatment in pediatric patients with cardiac surgery. *Clin Res Cardiol* 2008;97:801-10.
298. Adamson PC, Rhodes LA, Saul JP, et al. The pharmacokinetics of esmolol in pediatric subjects with supraventricular arrhythmias. *Pediatr Cardiol* 2006;27:420-7.
299. Chrysostomou C, Beerman L, Shiderly D, Berry D, Morell VO, Munoz R. Dexmedetomidine: a novel drug for the treatment of atrial and junctional tachyarrhythmias during the perioperative period for congenital cardiac surgery: a preliminary study. *Anesth Analg* 2008;107:1514-22.
300. Benson Jr D, Smith W, Dunnigan A, Sterba R, Gallagher J. Mechanisms of regular wide QRS tachycardia in infants and children. *Am J Cardiol* 1982;49: 1778-88.
301. Drago F, Mazza A, Guccione P, Mafriaci A, Di Liso G, Ragonese P. Amiodarone used alone or in combination with propranolol: a very effective therapy for tachyarrhythmias in infants and children. *Pediatr Cardiol* 1998;19:445-9.
302. Benson DJ, Dunnigan A, Green T, Benditt D, Schneider S. Periodic procainamide for paroxysmal tachycardia. *Circulation* 1985;72:147-52.
303. Komatsu C, Ishinaga T, Tateishi O, Tokuhisa Y, Yoshimura S. Effects of four antiarrhythmic drugs on the induction and termination of paroxysmal supraventricular tachycardia. *Jpn Circ J* 1986;50:961-72.
304. Mandel WJ, Laks MM, Obayashi K, Hayakawa H, Daley W. The WolffParkinsonWhite syndrome: pharmacologic effects of procaine amide. *Am Heart J* 1975;90:744-54.
305. Meldon SW, Brady WJ, Berger S, Mannenbach M. Pediatric ventricular tachycardia: a review with three illustrative cases. *Pediatr Emerg Care* 1994;10:294-300.

306. Shih JY, Gillette PC, Kugler JD, et al. The electrophysiologic effects of procainamide in the immature heart. *Pediatr Pharmacol (New York)* 1982;2:65–73.
307. Ackerman MJ, Siu BL, Sturner WQ, et al. Postmortem molecular analysis of SCN5A defects in sudden infant death syndrome. *JAMA* 2001;286:2264–9.
308. Arnestad M, Crotti L, Rognum TO, et al. Prevalence of longQT syndrome gene variants in sudden infant death syndrome. *Circulation* 2007;115: 361–7.
309. Cronk LB, Ye B, Kaku T, et al. Novel mechanism for sudden infant death syndrome: persistent late sodium current secondary to mutations in caveolin3. *Heart Rhythm* 2007;4:161–6.
310. Millat G, Kugener B, Chevalier P, et al. Contribution of longQT syndrome genetic variants in sudden infant death syndrome. *Pediatr Cardiol* 2009;30:502–9.
311. Otogiri T, Kijima K, Osawa M, et al. Cardiac ion channel gene mutations in sudden infant death syndrome. *Pediatr Res* 2008;64:482–7.
312. Plant LD, Bowers PN, Liu Q, et al. A common cardiac sodium channel variant associated with sudden infant death in African Americans, SCN5A S1103Y. *J Clin Invest* 2006;116:430–5.
313. Tester DJ, Dura M, Carturan E, et al. A mechanism for sudden infant death syndrome (SIDS): stress-induced leak via ryanodine receptors. *Heart Rhythm* 2007;4:733–9.
314. Albert CM, Nam EG, Rimm EB, et al. Cardiac sodium channel gene variants and sudden cardiac death in women. *Circulation* 2008;117:16–23.
315. Chugh SS, Senashova O, Watts A, et al. Postmortem molecular screening in unexplained sudden death. *J Am Coll Cardiol* 2004;43:1625–9.
316. Tester DJ, Spoon DB, Valdivia HH, Makielski JC, Ackerman MJ. Targeted mutational analysis of the RyR2-encoded cardiac ryanodine receptor in sudden unexplained death: a molecular autopsy of 49 medical examiner/coroner's cases. *Mayo Clin Proc* 2004;79:1380–4.
317. Calkins CM, Bensard DD, Partrick DA, Karrer FM. A critical analysis of outcome for children sustaining cardiac arrest after blunt trauma. *J Pediatr Surg* 2002;37:180–4.
318. Crewdson K, Lockey D, Davies G. Outcome from paediatric cardiac arrest associated with trauma. *Resuscitation* 2007;75:29–34.
319. Lopez-Herce Cid J, Dominguez Sampedro P, Rodriguez Nunez A, et al. Cardiorespiratory arrest in children with trauma. *An Pediatr (Barc)* 2006;65: 439–47.
320. Perron AD, Sing RF, Branas CC, Huynh T. Predicting survival in pediatric trauma patients receiving cardiopulmonary resuscitation in the prehospital setting. *Prehosp Emerg Care* 2001;5:6–9.
321. Sheikh A, Brogan T. Outcome and cost of open and closed chest cardiopulmonary resuscitation in pediatric cardiac arrests. *Pediatrics* 1994;93: 392–8.
322. Beaver BL, Colombani PM, Buck JR, Dudgeon DL, Bohrer SL, Haller Jr JA. Efficacy of emergency room thoracotomy in pediatric trauma. *J Pediatr Surg* 1987;22:19–23.
323. Powell RW, Gill EA, Jurkovich GJ, Ramenofsky ML. Resuscitative thoracotomy in children and adolescents. *Am Surg* 1988;54:188–91.
324. Rothenberg SS, Moore EE, Moore FA, Baxter BT, Moore JB, Cleveland HC. Emergency department thoracotomy in children – a critical analysis. *J Trauma* 1989;29:1322–5.
325. Suominen P, Rasanen J, Kivioja A. Efficacy of cardiopulmonary resuscitation in pulseless paediatric trauma patients. *Resuscitation* 1998;36:9–13.
326. Graham EM, Forbus GA, Bradley SM, Shirali GS, Atz AM. Incidence and outcome of cardiopulmonary resuscitation in patients with shunted single ventricle: advantage of right ventricle to pulmonary artery shunt. *J Thorac Cardiovasc Surg* 2006;131:e7–8.
327. Charpie JR, Dekeon MK, Goldberg CS, Mosca RS, Bove EL, Kulik TJ. Postoperative hemodynamics after Norwood palliation for hypoplastic left heart syndrome. *Am J Cardiol* 2001;87:198–202.
328. Hoffman GM, Mussatto KA, Brosig CL, et al. Systemic venous oxygen saturation after the Norwood procedure and childhood neurodevelopmental outcome. *J Thorac Cardiovasc Surg* 2005;130:1094–100.
329. Johnson BA, Hoffman GM, Tweddell JS, et al. Near-infrared spectroscopy in neonates before palliation of hypoplastic left heart syndrome. *Ann Thorac Surg* 2009;87:571–7, discussion 7–9.
330. Hoffman GM, Tweddell JS, Ghanayem NS, et al. Alteration of the critical arteriovenous oxygen saturation relationship by sustained afterload reduction after the Norwood procedure. *J Thorac Cardiovasc Surg* 2004;127:738–45.
331. De Oliveira NC, Van Arsdell GS. Practical use of alpha blockade strategy in the management of hypoplastic left heart syndrome following stage one palliation with a Blalock-Taussig shunt. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu* 2004;7:11–5.
332. Tweddell JS, Hoffman GM, Mussatto KA, et al. Improved survival of patients undergoing palliation of hypoplastic left heart syndrome: lessons learned from 115 consecutive patients. *Circulation* 2002;106:182–9.

333. Shekerdemian LS, Shore DF, Lincoln C, Bush A, Redington AN. Negative pressure ventilation improves cardiac output after right heart surgery. *Circulation* 1996;94:1149–55.
334. Shekerdemian LS, Bush A, Shore DF, Lincoln C, Redington AN. Cardiopulmonary interactions after Fontan operations: augmentation of cardiac output using negative pressure ventilation. *Circulation* 1997;96:3934–42.
335. Booth KL, Roth SJ, Thiagarajan RR, Almodovar MC, del Nido PJ, Laussen PC. Extracorporeal membrane oxygenation support of the Fontan and bidirectional Glenn circulations. *Ann Thorac Surg* 2004;77:1341–8.
336. Polderman FN, Cohen J, Blom NA, et al. Sudden unexpected death in children with a previously diagnosed cardiovascular disorder. *Int J Cardiol* 2004;95:171–6.
337. Sanatani S, Wilson G, Smith CR, Hamilton RM, Williams WG, Adatia I. Sudden unexpected death in children with heart disease. *Congenit Heart Dis* 2006;1:89–97.
338. Morris K, Beghetti M, Petros A, Adatia I, Bohn D. Comparison of hyperventilation and inhaled nitric oxide for pulmonary hypertension after repair of congenital heart disease. *Crit Care Med* 2000;28:2974–8.
339. Hoepfer MM, Galie N, Murali S, et al. Outcome after cardiopulmonary resuscitation in patients with pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2002;165:341–4.
340. Rimensberger PC, SpahrSchopfer I, Berner M, et al. Inhaled nitric oxide versus aerosolized iloprost in secondary pulmonary hypertension in children with congenital heart disease: vasodilator capacity and cellular mechanisms. *Circulation* 2001;103:544–8.
341. Liu KS, Tsai FC, Huang YK, et al. Extracorporeal life support: a simple and effective weapon for postcardiotomy right ventricular failure. *Artif Organs* 2009;33:504–8.
342. Dhillon R, Pearson GA, Firmin RK, Chan KC, Leanage R. Extracorporeal membrane oxygenation and the treatment of critical pulmonary hypertension in congenital heart disease. *Eur J Cardiothorac Surg* 1995;9:553–6.
343. Arpesella G, Loforte A, Mikus E, Mikus PM. Extracorporeal membrane oxygenation for primary allograft failure. *Transplant Proc* 2008;40:3596–7.
344. Strueber M, Hoepfer MM, Fischer S, et al. Bridge to thoracic organ transplantation in patients with pulmonary arterial hypertension using a pumpless lung assist device. *Am J Transplant* 2009;9:853–7.
345. Nolan JP, Neumar RW, Adrie C, et al. Postcardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A Scientific Statement from the International Liaison Committee on Resuscitation; the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; the Council on Stroke. *Resuscitation* 2008;79:350–79.
346. Hildebrand CA, Hartmann AG, Arcinue EL, Gomez RJ, Bing RJ. Cardiac performance in pediatric neardrowning. *Crit Care Med* 1988;16:331–5.
347. Checchia PA, Sehra R, Moynihan J, Daher N, Tang W, Weil MH. Myocardial injury in children following resuscitation after cardiac arrest. *Resuscitation* 2003;57:131–7.
348. Mayr V, Luckner G, Jochberger S, et al. Arginine vasopressin in advanced cardiovascular failure during the postresuscitation phase after cardiac arrest. *Resuscitation* 2007;72:35–44.
349. Huang L, Weil MH, Sun S, Cammarata G, Cao L, Tang W. Levosimendan improves postresuscitation outcomes in a rat model of CPR. *J Lab Clin Med* 2005;146:256–61.
350. Huang L, Weil MH, Tang W, Sun S, Wang J. Comparison between dobutamine and levosimendan for management of postresuscitation myocardial dysfunction. *Crit Care Med* 2005;33:487–91.
351. Kern KB, Hilwig RW, Rhee KH, Berg RA. Myocardial dysfunction after resuscitation from cardiac arrest: an example of global myocardial stunning. *J Am Coll Cardiol* 1996;28:232–40.
352. Meyer RJ, Kern KB, Berg RA, Hilwig RW, Ewy GA. Postresuscitation right ventricular dysfunction: delineation and treatment with dobutamine. *Resuscitation* 2002;55:187–91.
353. Studer W, Wu X, Siegemund M, Marsch S, Seeberger M, Filipovic M. Influence of dobutamine on the variables of systemic haemodynamics, metabolism, and intestinal perfusion after cardiopulmonary resuscitation in the rat. *Resuscitation* 2005;64:227–32.
354. Vasquez A, Kern KB, Hilwig RW, Heidenreich J, Berg RA, Ewy GA. Optimal dosing of dobutamine for treating postresuscitation left ventricular dysfunction. *Resuscitation* 2004;61:199–207.
355. Hoffman TM, Wernovsky G, Atz AM, et al. Efficacy and safety of milrinone in preventing low cardiac output syndrome in infants and children after corrective surgery for congenital heart disease. *Circulation* 2003;107:996–1002.
356. Alvarez J, Bouzada M, Fernandez AL, et al. Hemodynamic effects of levosimendan compared with dobutamine in patients with low cardiac output after cardiac surgery. *Rev Esp Cardiol* 2006;59:338–45.

357. Jorgensen K, BechHanssen O, Houltz E, Ricksten SE. Effects of levosimendan on left ventricular relaxation and early filling at maintained preload and afterload conditions after aortic valve replacement for aortic stenosis. *Circulation* 2008;117:1075–81.
358. Lobato EB, Willert JL, Looke TD, Thomas J, Urdaneta F. Effects of milrinone versus epinephrine on left ventricular relaxation after cardiopulmonary bypass following myocardial revascularization: assessment by color mmode and tissue Doppler. *J Cardiothorac Vasc Anesth* 2005;19:334–9.
359. Nijhawan N, Nicolosi AC, Montgomery MW, Aggarwal A, Pagel PS, Warltier DC. Levosimendan enhances cardiac performance after cardiopulmonary bypass: a prospective, randomized placebocontrolled trial. *J Cardiovasc Pharmacol* 1999;34:219–28.
360. Hickey RW, Kochanek PM, Ferimer H, Graham SH, Safar P. Hypothermia and hyperthermia in children after resuscitation from cardiac arrest. *Pediatrics* 2000;106:118–22.
361. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;346:549–56.
362. Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of outofhospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;346:557–63.
363. Gluckman PD, Wyatt JS, Azzopardi D, et al. Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicentre randomised trial. *Lancet* 2005;365:663–70.
364. Battin MR, Penrice J, Gunn TR, Gunn AJ. Treatment of term infants with head cooling and mild systemic hypothermia (35.0 degrees C and 34.5 degrees C) after perinatal asphyxia. *Pediatrics* 2003;111:244–51.
365. Compagnoni G, Pogliani L, Lista G, Castoldi F, Fontana P, Mosca F. Hypothermia reduces neurological damage in asphyxiated newborn infants. *Biol Neonate* 2002;82:222–7.
366. Gunn AJ, Gunn TR, Gunning MI, Williams CE, Gluckman PD. Neuroprotection with prolonged head cooling started before postischemic seizures in fetal sheep. *Pediatrics* 1998;102:1098–106.
367. Debillon T, Daoud P, Durand P, et al. Wholebody cooling after perinatal asphyxia: a pilot study in term neonates. *Dev Med Child Neurol* 2003;45: 17–23.
368. Shankaran S, Laptook AR, Ehrenkranz RA, et al. Wholebody hypothermia for neonates with hypoxicischemic encephalopathy. *N Engl J Med* 2005;353:1574–84.
369. Doherty DR, Parshuram CS, Gaboury I, et al. Hypothermia therapy after pediatric cardiac arrest. *Circulation* 2009;119:1492–500.
370. HachimiIdrissi S, Corne L, Ebinger G, Michotte Y, Huyghens L. Mild hypothermia induced by a helmet device: a clinical feasibility study. *Resuscitation* 2001;51:275–81.
371. Bernard S, Buist M, Monteiro O, Smith K. Induced hypothermia using large volume, icecold intravenous fluid in comatose survivors of outofhospital cardiac arrest: a preliminary report. *Resuscitation* 2003;56:9–13.
372. Kliegel A, Losert H, Sterz F, et al. Cold simple intravenous infusions preceding special endovascular cooling for faster induction of mild hypothermia after cardiac arrest – a feasibility study. *Resuscitation* 2005;64:347–51.
373. Polderman KH. Application of therapeutic hypothermia in the ICU: opportunities and pitfalls of a promising treatment modality. Part 1: Indications and evidence. *Intensive Care Med* 2004;30:556–75.
374. Polderman KH. Application of therapeutic hypothermia in the intensive care unit. Opportunities and pitfalls of a promising treatment modality. Part 2: Practical aspects and side effects. *Intensive Care Med* 2004;30:757–69.
375. Polderman KH, Peerdeman SM, Girbes AR. Hypophosphatemia and hypomagnesemia induced by cooling in patients with severe head injury. *J Neurosurg* 2001;94:697–705.
376. Zeiner A, Holzer M, Sterz F, et al. Hyperthermia after cardiac arrest is associated with an unfavorable neurologic outcome. *Arch Intern Med* 2001;161:2007–12.
377. Takino M, Okada Y. Hyperthermia following cardiopulmonary resuscitation. *Intensive Care Med* 1991;17:419–20.
378. Takasu A, Saitoh D, Kaneko N, Sakamoto T, Okada Y. Hyperthermia: is it an ominous sign after cardiac arrest? *Resuscitation* 2001;49:273–7.
379. Coimbra C, BorisMoller F, Drake M, Wieloch T. Diminished neuronal damage in the rat brain by late treatment with the antipyretic drug dipyron or cooling following cerebral ischemia. *Acta Neuropathol* 1996;92:447–53.
380. Coimbra C, Drake M, BorisMoller F, Wieloch T. Longlasting neuroprotective effect of postischemic hypothermia and treatment with an antiinflammatory/ antipyretic drug. Evidence for chronic encephalopathic processes following ischemia. *Stroke* 1996;27:1578–85.
381. van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in the critically ill patients. *N Engl J Med* 2001;345:1359–67.

382. Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med* 2006;354:449–61.
383. Wiener RS, Wiener DC, Larson RJ. Benefits and risks of tight glucose control in critically ill adults: a metaanalysis. *JAMA* 2008;300:933–44.
384. TreggiariMM, Karir V, Yanez ND, Weiss NS, Daniel S, Deem SA. Intensive insulin therapy and mortality in critically ill patients. *Crit Care* 2008;12:R29.
385. Slonim AD, Patel KM, Ruttimann UE, Pollack MM. Cardiopulmonary resuscitation in pediatric intensive care units. *Crit Care Med* 1997;25:1951–5.
386. RodriguezNunez A, LopezHerce J, Garcia C, et al. Effectiveness and longterm outcome of cardiopulmonary resuscitation in paediatric intensive care units in Spain. *Resuscitation* 2006;71:301–9.
387. Nadkarni VM, Larkin GL, Peberdy MA, et al. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. *JAMA* 2006;295:50–7.
388. Meaney PA, Nadkarni VM, Cook EF, et al. Higher survival rates among younger patients after pediatric intensive care unit cardiac arrests. *Pediatrics* 2006;118:2424–33.
389. Tibballs J, Kinney S. A prospective study of outcome of inpatient paediatric cardiopulmonary arrest. *Resuscitation* 2006;71:310–8.
390. Gillis J, Dickson D, Rieder M, Steward D, Edmonds J. Results of inpatient pediatric resuscitation. *Crit Care Med* 1986;14:469–71.
391. Schindler MB, Bohn D, Cox PN, et al. Outcome of outofhospital cardiac or respiratory arrest in children. *N Engl J Med* 1996;335:1473–9.
392. Suominen P, Korpela R, Kuisma M, Silfvast T, Olkkola KT. Paediatric cardiac arrest and resuscitation provided by physicianstaffed emergency care units. *Acta Anaesthesiol Scand* 1997;41:260–5.
393. LopezHerce J, Garcia C, Dominguez P, et al. Outcome of outofhospital cardiorespiratory arrest in children. *Pediatr Emerg Care* 2005;21:807–15.
394. LopezHerce J, Garcia C, Dominguez P, et al. Characteristics and outcome of cardiorespiratory arrest in children. *Resuscitation* 2004;63:311–20.
395. Hazinski MF, Chahine AA, Holcomb 3rd GW, Morris Jr JA. Outcome of cardiovascular collapse in pediatric blunt trauma. *Ann Emerg Med* 1994;23:1229–35.
396. MorrisMC, Wernovsky G, NadkarniVM. Survival outcomes after extracorporeal cardiopulmonary resuscitation instituted during active chest compressions following refractory in-hospital pediatric cardiac arrest. *Pediatr Crit Care Med* 2004;5:440–6.
397. Duncan BW, Ibrahim AE, Hraska V, et al. Use of rapiddeployment extracorporeal membrane oxygenation for the resuscitation of pediatric patients with heart disease after cardiac arrest. *J Thorac Cardiovasc Surg* 1998;116: 305–11.
398. Parra DA, Totapally BR, Zahn E, et al. Outcome of cardiopulmonary resuscitation in a pediatric cardiac intensive care unit. *Crit Care Med* 2000;28: 3296–300.
399. Idris AH, Berg RA, Bierens J, et al. Recommended guidelines for uniform reporting of data from drowning: the “Utstein style”. *Resuscitation* 2003;59:45–57.
400. Eich C, Brauer A, Timmermann A, et al. Outcome of 12 drowned children with attempted resuscitation on cardiopulmonary bypass: an analysis of variables based on the “Utstein Style for Drowning”. *Resuscitation* 2007;75:42–52.
401. Dudley NC, Hansen KW, Furnival RA, Donaldson AE, Van Wagenen KL, Scaife ER. The effect of family presence on the efficiency of pediatric trauma resuscitations. *Ann Emerg Med* 2009;53:777–84, e3.
402. Tinsley C, Hill JB, Shah J, et al. Experience of families during cardiopulmonary resuscitation in a pediatric intensive care unit. *Pediatrics* 2008;122: e799–804.
403. Mangurten J, Scott SH, Guzzetta CE, et al. Effects of family presence during resuscitation and invasive procedures in a pediatric emergency department. *J Emerg Nurs* 2006;32:225–33.
404. McGaheyOakland PR, Lieder HS, Young A, et al. Family experiences during resuscitation at a children’s hospital emergency department. *J Pediatr Health Care* 2007;21:217–25.
405. Jones M, Qazi M, Young KD. Ethnic differences in parent preference to be present for painful medical procedures. *Pediatrics* 2005;116:e191–7.
406. Boie ET, Moore GP, Brummett C, Nelson DR. Do parents want to be present during invasive procedures performed on their children in the emergency department? A survey of 400 parents. *Ann Emerg Med* 1999;34: 70–4.
407. Andrews R, Andrews R. Family presence during a failed major trauma resuscitation attempt of a 15yearold boy: lessons learned [see comment]. *J Emerg Nurs* 2004;30:556–8.
408. Dill K, GanceCleveland B, Dill K, GanceCleveland B. With you until the end: family presence during failed resuscitation. *J Specialists Pediatr Nurs: JSPN* 2005;10:204–7.

409. Gold KJ, Gorenflo DW, Schwenk TL, et al. Physician experience with family presence during cardiopulmonary resuscitation in children [see comment]. *Pediatr Crit Care Med* 2006;7:428–33.
410. Duran CR, Oman KS, Abel JJ, Koziel VM, Szymanski D. Attitudes toward and beliefs about family presence: a survey of healthcare providers, patients' families, and patients. *Am J Crit Care* 2007;16:270–9.
411. Meyers TA, Eichhorn DJ, Guzzetta CE, et al. Family presence during invasive procedures and resuscitation. *Am J Nurs* 2000;100:32–42, quiz 3.
412. O'Connell KJ, Farah MM, Spandorfer P, et al. Family presence during pediatric trauma team activation: an assessment of a structured program. *Pediatrics* 2007;120:e565–74.
413. Engel KG, Barnosky AR, Berry B, et al. Provider experience and attitudes toward family presence during resuscitation procedures. *J Palliative Med* 2007;10:1007–9.
414. Holzhauser K, Finucane J, De Vries S. Family presence during resuscitation: a randomised controlled trial of the impact of family presence. *Australasian Emerg Nurs J* 2005;8:139–47.
415. Doyle CJ, Post H, Burney RE, Maino J, Keefe M, Rhee KJ. Family participation during resuscitation: an option. *Ann Emerg Med* 1987;16:673–5.
416. Meyers TA, Eichhorn DJ, Guzzetta CE. Do families want to be present during CPR? A retrospective survey. *J Emerg Nurs* 1998;24:400–5.
417. Hanson C, Strawser D. Family presence during cardiopulmonary resuscitation: foote Hospital emergency department's nine year perspective. *J Emerg Nurs* 1992;18:104–6.
418. Robinson SM, Mackenzie Ross S, Campbell Hewson GL, Egleston CV, Prevost AT. Psychological effect of witnessed resuscitation on bereaved relatives. *Lancet* 1998;352:614–7.
419. Compton S, Madgy A, Goldstein M, et al. Emergency medical service providers' experience with family presence during cardiopulmonary resuscitation. *Resuscitation* 2006;70:223–8.
420. Beckman AW, Sloan BK, Moore GP, et al. Should parents be present during emergency department procedures on children, and who should make that decision? A survey of emergency physician and nurse attitudes. *Acad Emerg Med* 2002;9:154–8.
421. Eppich WJ, Arnold LD. Family member presence in the pediatric emergency department. *Curr Opin Pediatr* 2003;15:294–8.
422. Eichhorn DJ, Meyers TA, Mitchell TG, Guzzetta CE. Opening the doors: family presence during resuscitation. *J Cardiovasc Nurs* 1996;10:59–70.
423. Jarvis AS. Parental presence during resuscitation: attitudes of staff on a paediatric intensive care unit. *Intensive Crit Care Nurs* 1998;14:3–7